



Potassium Channel Activation: A Potential Therapeutic Approach?

Kim Lawson

DIVISION OF BIOMEDICAL SCIENCES, SCHOOL OF SCIENCE,
SHEFFIELD HALLAM UNIVERSITY, CITY CAMPUS, POINTE STREET, SHEFFIELD S1 1WB, UK

ABSTRACT. The physiological role of K^+ channel opening by endogenous substances (e.g., neurotransmitters and hormones) is a recognised inhibitory mechanism. Thus, the identification of novel synthetic molecules that 'directly' open K^+ channels has led to a new direction in the pharmacology of ion channels. The existence of many different subtypes of K^+ channels has been an impetus in the search for new molecules demonstrating channel and, thus, tissue selectivity. This review focuses on the different classes of openers of K^+ channels, the intracellular mechanisms involved in the execution of their effects, and potential therapeutic targets. *PHARMACOL. THER.* 70(1): 39-63, 1996.

KEY WORDS. Potassium channel openers, KCO classification, K_{ATP} , K_{Ca} , cromakalim, therapeutic targets.

CONTENTS

| | | | |
|--------------------------------------|----|-----------------------------|----|
| 1. INTRODUCTION | 39 | 4.1.1. VASCULAR | 46 |
| 2. POTASSIUM CHANNEL OPENER | | 4.1.2. CARDIAC | 48 |
| CLASSIFICATION | 40 | 4.1.3. BLOOD | 48 |
| 2.1. ATP-SENSITIVE K^+ CHANNEL | | 4.2. RESPIRATORY SYSTEM | 48 |
| OPENERS | 41 | 4.3. REPRODUCTIVE SYSTEM | 50 |
| 2.2. CALCIUM-ACTIVATED POTASSIUM | | 4.4. URINARY BLADDER | 51 |
| CHANNEL OPENERS | 43 | 4.5. GASTROINTESTINAL TRACT | 51 |
| 2.3. ENDOGENOUS OPENERS | 43 | 4.6. NERVOUS SYSTEM | 52 |
| 2.4. OTHERS | 45 | 4.6.1. PERIPHERAL | 52 |
| 3. MECHANISMS OF ACTION | 45 | 4.6.2. CENTRAL | 52 |
| 3.1. HYPERPOLARIZATION | 45 | 4.7. SKELETAL MUSCLE | 53 |
| 3.2. INTRACELLULAR Ca^{2+} STORES | 45 | 4.8. HAIR GROWTH | 54 |
| 3.3. INOSITOL PHOSPHATE CASCADE | 46 | 4.9. INTRACULAR PRESSURE | 54 |
| 3.4. Ca^{2+} SENSITIVITY | 46 | 5. CONCLUSIONS | 54 |
| 3.5. OTHERS | 46 | ACKNOWLEDGEMENTS | 55 |
| 4. THERAPEUTIC TARGETS AND POTENTIAL | | REFERENCES | 55 |
| 4.1. CARDIOVASCULAR SYSTEM | 46 | | |

ABBREVIATIONS. BK_{Ca} , high-conductance calcium-activated potassium channel; $[Ca^{2+}]_i$, intracellular Ca^{2+} concentration; cAMP, cyclic AMP; cGMP, cyclic GMP; CGRP, calcitonin gene-related peptide; DHS-4, dehydroxypasiponin 1; DMPP, 1,1-dimethyl-4-phenylpiperazinium; EDHF, endothelium-derived hyperpolarizing factor; EDHF, endothelium-derived relaxing factor; E_K , potassium equilibrium potential; ET-1, endothelin-1; 5-HT, 5-hydroxytryptamine; K_{ATP} , ATP-sensitive potassium channel; K_{Ca} , calcium-activated potassium channel; $K_{Ca}CO$, calcium-activated potassium channel opener; KCO, potassium channel opener; $K_{ATP}CO$, ATP-sensitive potassium channel opener; NANCG, nonadrenergic, noncholinergic excitatory neurons; NDP, nucleotide diphosphate; NO, nitric oxide; PGI_2 , prostacyclin; pS, conductance; SK_{Ca} , small-conductance calcium-activated potassium channel; SMC, smooth muscle cells; VOC , voltage-operated Ca^{2+} channel.

1. INTRODUCTION

The biological cell is an integral structure that responds to chemical and physical extracellular signals on its membrane, which are communicated to the intracellular processes through a variety of pathways. The transmembrane movement (i.e., efflux and influx) of ions (e.g., Ca^{2+} , Na^+ , K^+ , Cl^-) through plasmalemma channels are universal mechanisms used to execute or modulate physiological functions in living cells.

Potassium (K^+) specific channels are a diverse and ubiquitous group of ion channels and, thus, play a fundamental role in the modulation of cell excitability (Hille, 1984; Rudy, 1988). In the resting state of excitable or nonexcitable cells,

the concentration of K^+ outside the membrane (3-5 mM) is at least 25-fold lower than the K^+ concentration in the intracellular fluid (130-160 mM). Consequently, an outward current due to efflux of positively charged ions is generated by the opening of K^+ channels. The efflux of K^+ is a mechanism for recovering (repolarization), maintaining (clamping), and/or enhancing (hyperpolarization) the resting potential of the cell. Thus, the opening of K^+ channels is a physiological means for counteracting, restricting, or preventing depolarizing activity caused by inward currents, due to entry of Ca^{2+} and Na^+ and the efflux of Cl^- ions.

The functions of these channels, which are crucial for subserving different physiological functions, depend on the

specific manners in which a particular K^+ channel opens and closes, and its selective permeation by K^+ ions. K^+ channels are generally classified according to their primary regulatory or gating mechanism (Hille, 1984). Ion channels can be characterised by ionic selectivity (differential permeability), conductance (pS), gating properties (factors controlling channel opening and closing), kinetics (rates at which channels open and close), and pharmacology (action of specific agents in blocking or changing the flow of ions). K^+ channel classifications and their pharmacological properties have been reviewed extensively (Rudy, 1988; Castle *et al.*, 1989; Cook, 1990; Dreyer, 1990; Kolb, 1990; Robertson and Steinberg, 1990; Atkinson, 1992). This article, however, will focus on the subtypes of K^+ channels (e.g., ATP-sensitive potassium channel [K_{ATP}], calcium-activated potassium channel [K_{Ca}]) that have been associated with synthetic and endogenous openers of these channels.

The discovery and development of selective ligands that interact with specific K^+ channels, together with the combination of recent electrophysiological and molecular biology techniques, has resulted in a more detailed characterization of the role these channels play in regulating cell function. An appreciation of the primary amino acid sequence of each K^+ channel protein, through molecular biology techniques, will allow the development of highly selective ligands that open a designated channel subtype. Such ligands would provide essential information regarding the physiological and pathophysiological importance of particular K^+ channels; thus, leading to the identification of drugs focused for defined clinical conditions.

The main objective of this review is to present the different classes of agents that open K^+ channels and discuss potential clinical targets.

2. POTASSIUM CHANNEL OPENER CLASSIFICATION

The emergence of synthetic potassium channel openers (KCOs) has only occurred recently; however, the physiological role of K^+ channel opening by endogenous substances (neurotransmitters and hormones) is a recognized inhibitory mechanism (Kurachi *et al.*, 1986; Bulbring and Tamita, 1987). The term "potassium channel openers" was introduced to describe a group of novel synthetic molecules, typified by cromakalim (Hamilton *et al.*, 1986; Hamilton and Weston, 1989), that have led to a new direction in the pharmacology of ion channels. Hamilton *et al.* (1986), reporting that cromakalim evoked smooth muscle relaxant effects by the opening of K^+ channels in cell membranes, initiated major research efforts in the search for other such molecules and in the determination of the specific channels involved. KCO properties have, subsequently, been demonstrated in a diverse range of synthetic chemical structures and endogenous substances. In addition, the existence of so many different subtypes of K^+ channel has been an impetus in the search for new molecules that would have profiles and channel selectivities different from those exhibited by the group of KCOs typified by cromakalim. Recent progress has been reported

in the search for agents that 'selectively' open calcium-activated K^+ channels (see Section 2.2). In addition to the development of synthetic molecules, a number of endogenous substances have been identified that exert some or all of their effects via K^+ channel opening (see Section 2.3).

The recent advances made in this area of research have brought into question the use of the broad term "potassium channel openers" for reference to a limited group of molecules (i.e., K_{ATP} openers, those typified by cromakalim). For the purpose of this review, the term "potassium channel opener" (and the abbreviation KCO) will be used in reference to any substance reported to open K^+ channels and, thus, associated with an efflux of K^+ ions from the cell. A subscript to the K of the abbreviation KCO will be used to indicate the postulated channel involved (e.g., K_{ATP} CO for openers of K_{ATP} channels).

Molecules (primarily K_{ATP} COs) exhibiting properties compatible with the opening of K^+ channels have been essential tools in the development of research approaches and pharmacological techniques within this area. Such techniques not only have assisted in the identification of other KCOs, but are crucial in the determination of the mechanisms of action and K^+ channel subtype involved.

The unidirectional movements of specific ions can be assessed by isotope flux techniques. The efflux of K^+ from a cell can be measured by using $^{42}K^+$ or the more stable tracer $^{86}Rb^+$. Briefly, the tissue or cells are loaded with the isotope during an incubation period (Quast and Baumlin, 1988). The preparation is then challenged with a KCO and the radioactivity within the bathing environment and the residual radioactivity in the tissue is determined, thereby giving qualitative and quantitative representation of K^+ ion movements. In experimental models where mechanical responses can be recorded, the K^+ flux can be compared with functional responses following exposure of the tissue to the KCO.

KCOs hyperpolarize the cell membrane up to values approaching the potassium equilibrium potential (E_K ; see Section 3.1). Electrophysiological techniques permit the recording of transmembrane ion movements and membrane potential, depending on the experimental methodology, at a high level of resolution. Patch-clamp procedure applied to whole-cell configuration allows macroscopic K^+ currents to be recorded when depolarizing pulses are delivered to the preparation clamped at negative potentials close to E_K (Hamill *et al.*, 1981). Recording from isolated membrane patches also provides measurements of electrical currents through single K^+ channels (Hamill *et al.*, 1981). The difficulty in identifying certain K^+ channel subtypes in some tissues using these techniques may be due to their low density, fragility to the isolation procedure, and/or their irreversible rundown (where the activity of the K^+ channels recorded in cell free patches spontaneously declines over a relatively short period in the absence of the cytosol). Thus, experimental conditions, such as the isolation procedure and recording techniques, can influence the determination of the properties of K^+ channels.

Radioiodine binding assays have identified binding sites

for selected $K_{ATP}COs$, [PHIP1075 (pinacidil analog) and [Hromakalim], in smooth muscle preparations (Howlett and Longman, 1992; Quast *et al.*, 1993). [PHIP1075 binding in *rat* aorta was inhibited by representatives from all chemical families of $K_{ATP}COs$ with potencies that correlated with the potencies obtained in $^{86}Rb^+$ efflux and vasorelaxation studies (Quast *et al.*, 1993). Differences in the data obtained with [PHIP1075 and [Hromakalim] in smooth muscle preparations, however, suggest the existence of two different binding sites for $K_{ATP}COs$ (Lawson and Hicks, 1993). The use of strips of tissue with intact cells, not membrane preparations, being critical in this assay would suggest that the binding of these $K_{ATP}COs$ is dependent on the functional integrity of the cell (Quast *et al.*, 1993). The development of similar assays for other subtypes of K^+ channel, however, are dependent on the availability of appropriate ligands. Ligand binding studies on expressed cloned K^+ channels (e.g., ROMK1; Ho *et al.*, 1993), which should provide major advances in our understanding of how $KCOs$ interact with the channel, are awaited.

Functional isolated organ preparations have been used extensively in KCO research to determine the mechanical effects due to exposure of tissues to such compounds. Studies in parallel with those techniques described above can demonstrate whether or not the functional responses to $KCOs$ are compatible with and, thereby, a consequence of, K^+ channel opening and K^+ flux. Finally, the effects of KCO are determined in preclinical *in vivo* models predictive of potential therapeutic applications (see Section 4).

2.1. ATP-Sensitive K^+ Channel Openers

K_{ATP} channels, which have been studied extensively, initially were identified in cardiac cells and pancreatic β -cells (Noma, 1983; Ashcroft and Ashcroft, 1990). The intracellular concentration of ATP determines the state (open, closed) of the K_{ATP} channel (Edwards and Weston, 1993). Conk and Males (1984) proposed that K_{ATP} channels in pancreatic β -cells are spontaneously open under normal conditions and, as a result, a basal efflux of K^+ from the cell leads to hyperpolarization of the cell membrane. An increase in the glucose levels evokes a rise in intracellular ATP levels, closing the K_{ATP} channel. The resultant membrane depolarization and subsequent Ca^{2+} influx through voltage-operated Ca^{2+} channels ($VOCs$) stimulates insulin release. Exogenous compounds, in particular the sulphonylureas (e.g., glibenclamide), by closing K_{ATP} channels, can induce the release of insulin from pancreatic β -cells.

Subsequent studies have shown K_{ATP} channels to exist in virtually all tissues studied, including skeletal muscle, smooth muscle and neuronal cells (Ashcroft and Ashcroft, 1990). Five different types of K_{ATP} channels have been defined as a consequence of potassium selectivity and sensitivity to calcium, intracellular ATP concentration, and pharmacological modulation (Ashcroft and Ashcroft, 1990). Full pharmacological characterization of the putative subtypes of K_{ATP} channels is presently limited (Ashcroft and Ashcroft, 1990; Gopikrishnan *et al.*, 1993). Of these channels,

the Type I, which are blocked by micromolar concentrations of intracellular ATP, have been extensively studied in a number of cell types. This channel, irrespective of location, always demonstrates sensitivity to sulphonylureas; however, the concentrations of these compounds required to block the K_{ATP} channels are tissue-dependent (Edwards and Weston, 1993). In pancreatic β -cells, modulation of insulin release occurs at nanomolar concentrations of glibenclamide, and micromolar concentrations are required in smooth muscle and cardiac preparations, suggesting at least two concentration ranges of activity. Further, the rank order of potency of $K_{ATP}COs$ on Type I K_{ATP} channels is dependent on the cell type being studied (Edwards and Weston, 1993) and, thus, is open to further differentiation on the basis of pharmacological profile. In addition, the current nomenclature does not take into account the results of recent studies (albeit limited) using molecular biology of the cloned K_{ATP} channel (Ho *et al.*, 1993).

Classification of compounds termed $K_{ATP}COs$ has largely been defined as a consequence of their biological effects being sensitive to blockade by sulphonylureas. Potassium channel opening properties, as identified by electrophysiological and efflux studies, sensitive to glibenclamide (or other sulphonylureas) have been demonstrated in a diverse range of synthetic chemical structures (benzothiadiazines [e.g., diazoxide], pyrimidines [e.g., minoxidil], pyridylcyanoguanidines [e.g., pinacidil], nicotinamides [e.g., nicotidil], benzopyrans [e.g., cromakalim] and carbobenzimidazoles [e.g., RP 49356]) (Edwards and Weston, 1990; Lawson and Hicks, 1993). Common chemical structural features between benzopyrans, pyridylcyanoguanidines, and carbobenzimidazoles have been described (Arwal, 1992). In addition, most of the biological activity resides predominantly in the (-)-enantiomers of cromakalim, pinacidil, and RP 49356.

In functional *in vitro* pharmacology studies, the antagonism by glibenclamide of the effects of $K_{ATP}COs$ on vascular (Elrre, 1989a; Wilson, 1989; Newgreen *et al.*, 1990) and certain nonvascular (Elrre, 1989b; Pipet *et al.*, 1990; Edwards *et al.*, 1991) smooth muscle preparations has been described as competitive in nature. Competitive antagonism is inferred by parallel displacements of the $K_{ATP}CO$ -induced relaxant concentration-response curves to the right of controls, without a reduction of the maximum effect to the agonist by the antagonist, and Schild analysis (Arunkrishnan and Schild, 1959) yielding a slope not different from unity. The failure of glibenclamide (up to 10 μM) to displace [PHromakalim from its binding sites on rat aorta (Howlett and Longman, 1992) does not support a competitive interaction between these two compounds at a single site. Glibenclamide concentration dependently increased the dissociation rate of the [PHIP1075 binding complex on rat aorta (Quast *et al.*, 1993), suggesting that the glibenclamide site is distinct from, but negative allosterically coupled to, the binding site for the openers.

Parallel displacement of concentration-response curves could also be observed with physiological antagonism and, also, if spare channels were available (i.e., the $K_{ATP}CO$ can

evokes a maximal response without the activation of all of the channels in the tissue. The smooth muscle relaxant responses to K_{ATP} COs in the guinea-pig trachea, however, are blocked by glibenclamide in a manner that is not consistent with competitive antagonism (i.e. the maximum effect of the K_{ATP} CO is reduced in the presence of the antagonist) (Nielsen-Kudsk *et al.*, 1990; Berry *et al.*, 1991; Small *et al.*, 1992), but is consistent with a lack of spare channels (i.e. the K_{ATP} CO must activate all channels in the tissue to evoke a maximal response). Thus, glibenclamide appears to interact differently with the channel activated by K_{ATP} COs in respiratory smooth muscle from that in vascular tissue.

In vascular preparations, glibenclamide antagonized the effects of cromakalim competitively, whereas the effects of minoxidil sulphate were antagonized in a noncompetitive manner (Wickenden *et al.*, 1991). These findings led to the suggestion of the existence of different subtypes of K^+ channel sensitive to K_{ATP} COs. In addition, minoxidil sulphate, in contrast to dioxazine and cromakalim, had no effect on $^{86}Rb^+$ efflux in rat aorta, but did increase $^{42}K^+$ efflux (Newgreen *et al.*, 1990; Wickenden *et al.*, 1991). All three K_{ATP} COs increased $^{86}Rb^+$ and $^{42}K^+$ efflux from rat portal vein (Newgreen *et al.*, 1990). Thus, in rat aorta, minoxidil sulphate may open a K^+ channel impermeable to $^{86}Rb^+$ and not recognised by other K_{ATP} COs.

A number of anomalies exist within the pharmacology of the different chemical structures that have been tentatively classified into the family of K_{ATP} openers (Lawson and Hicks, 1993). A common feature responsible for the grouping of these compounds has been the susceptibility of the biological effects to inhibition by sulphonylureas. Nevertheless, the rank order of potency of the inhibitory activity and nature of antagonism by the sulphonylureas is dependent on the cell type and K_{ATP} CO being studied (Longman and Hamilton, 1992; Edwards and Weston, 1993; Lawson and Hicks, 1993). The pharmacological profile of (certain) K_{ATP} COs on the K_{ATP} channel is also dependent upon cell type (e.g., dioxazine is an 'activator' in pancreatic cells, inactive in skeletal muscle cells, and an 'antagonist' in cardiac cells) (Zankler *et al.*, 1988; Faivre and Findlay, 1989; Weik and Neumcke, 1990). Discrimination between the pharmacology of benzopyran and nonbenzopyran K_{ATP} COs has been identified (Lawson *et al.*, 1992; Randall and Griffith, 1993), where all compounds have exhibited definable KCO profiles. Such findings support the concept that there are differences in the way these agents interact with potassium channels, whichever subtype they may be (Lawson *et al.*, 1992; Lawson and Hicks, 1993).

The ability of glibenclamide to antagonize the actions of this group of compounds led to the suggestion of K_{ATP} channels being the site of action; however, this is still an area of research and discussion. Determination of a common site of action for this group of compounds is complicated by, at least, (1) the existence of subtypes of K_{ATP} channels, (2) evidence of these compounds opening several K^+ channels, and (3) effects of glibenclamide on K^+ channels other than K_{ATP} channels.

Using patch-clamp techniques on the rabbit portal vein, Beech *et al.* (1993a,b) found that nucleotide diphosphates (NDPs), such as GTP, are required for K_{ATP} channel opening in the absence of the inhibitory effects of ATP. This property sets them apart from the K_{ATP} channel present in the pancreas and heart. These authors suggested that this K^+ channel in rabbit portal vein should be more correctly termed K_{NDP} to reflect the obligatory role that NDPs play in regulating this particular channel. Together with the recent findings of Kitamura and Kamouchi (1993) that K_{ATP} channels appear to be nonuniform across species in various smooth muscle cells (SMC), evidence suggests that there may be several different types of ATP, adenosine nucleotide, and glibenclamide-sensitive K_{ATP} channels in various tissues that may be important targets for compounds that can selectively regulate their activity. These findings highlight the complexity of the biological system and the need to fully investigate the specificity of channel/ligand interactions to clearly define the site of action of openers of K_{ATP} channels.

A number of K_{ATP} COs, however, (e.g., pinacidil and nicorandil) possess additional properties (e.g., adenylylate or guanylate cyclase stimulation) to that of potassium channel opening that could account, in part, for pharmacological profiles differing from those of the benzopyrans (e.g., cromakalim) and carboxathiadines (e.g., RP 493561 (Cook, 1990; Longman and Hamilton, 1992). The situation is further complicated by agents defined as K_{ATP} COs and glibenclamide exhibiting the actions on K^+ channels other than K_{ATP} channels, Ca^{2+} channels and Cl^- currents. Glibenclamide can inhibit delayed rectifier K^+ currents (Reuve *et al.*, 1992; Beech *et al.*, 1993b; Crepel *et al.*, 1993). A type K^+ current (Sadraei and Beech, 1995), Ca^{2+} channel currents (Blum and Herrmeyer, 1994; Sadraei and Beech, 1995) and Cl^- current (Sheppard and Welsh, 1992). Cromakalim activated K_{Ca} current (Gelband and McCullough, 1993) and inhibited Ca^{2+} channel current (Okabe *et al.*, 1990) and Cystic Fibrosis Transmembrane Regulator-dependent Cl^- current (Sheppard and Welsh, 1992).

A heterogeneity in the pharmacology of the K_{ATP} COs can be identified, which would suggest a heterogeneity of the K_{ATP} sites of action and/or the K^+ channel. Interpretation of data is further complicated by the potential heterogeneity of sulphonylurea-sensitive sites of deactivation that may be independent of the K_{ATP} CO site of action. Further work involving comparative studies with representatives of each chemical family of K_{ATP} COs will assist in the understanding and subclassification of these heterogeneous sites of action. A combined effort of electrophysiology, functional pharmacology, and molecular biology, with a subsequent study of the expressed channels, will be required to address the real question of channel selectivity and K_{ATP} CO subclassification. The differences already observed in the pharmacology of K_{ATP} COs are, therefore, important factors to consider in the development of second generation compounds, where tissue or organ selectivity is sought.

2.2. Calcium-Activated Potassium Channel Openers

K_{Ca} channels are activated by membrane depolarization and by increases in intracellular calcium [Barr and Magleby, 1987]. Three subtypes of K_{Ca} channels have been described on the basis of their single-channel conductance and sensitivity to specific pharmacological blockers [Cook, 1990]. High-conductance or maxi-K (BK_{Ca} , 100–250 pS) channels are sensitive to charyldotoxin and ibetoxin, intermediate-conductance (IK_{Ca} , 18–50 pS) channels are blocked by high concentrations of charyldotoxin, and low- or small-conductance (SK_{Ca} , 10–16 pS) channels are potentially blocked by apamin. K_{Ca} channels have been identified in virtually all types of cells, where they almost certainly function to terminate excitatory processes that are triggered or maintained by an increase in the intracellular Ca^{2+} concentration ($[Ca^{2+}]_i$) and/or involve depolarization. As a consequence of such an important physiological role of K_{Ca} channels, the search and recent identification of selective synthetic BK_{Ca} openers has received significant attention.

Two major chemical groups, benzimidazoles (e.g., NS-004, NS-1619) and imidazo[2,1-*a*]pyridines (e.g., SCA40), recently have been reported to exhibit BK_{Ca} channel opening properties [Sahlayrolles et al., 1988; Olesen and Watjen, 1992; Laurent et al., 1993; Olesen et al., 1993]. NS-004 directly activates the charyldotoxin-sensitive BK_{Ca} channel in rat cerebellar granule cells, in rat (OH) clonal pituitary tumor cells, bovine aortic SMC and guinea-pig tracheal SMC [Olesen and Watjen, 1992; Olesen et al., 1993, 1994a; McKay et al., 1994]. NS-1619 directly activated the BK_{Ca} channel in rat ventromedial hypothalamic neurons, bovine aortic SMC, and rat portal vein [Edwards et al., 1994; Sellers and Ashford, 1994; Olesen et al., 1994b]. Although NS-1619 failed to modify the K_{ATP} channel in rat ventromedial hypothalamus [Sellers and Ashford, 1994], in the rat portal vein, this compound was reported to inhibit the KV channel over the same concentration range that activated the BK_{Ca} channel [Edwards et al., 1994]. NS-004 induced relaxation in guinea-pig trachea, which were sensitive to charyldotoxin and ibetoxin, but not glibenclamide (K_{ATP} channel), dolenilide (inward rectifier) or apamin (SK_{Ca}), suggesting activation of only BK_{Ca} channels being responsible for this functional response [Lawson et al., 1996]. Although NS-004 and NS-1619 activate BK_{Ca} channels, their relaxant effects in rat portal vein and cardioprotective effects probably are due to inhibition of L-type calcium channels [Edwards et al., 1994; Sargent et al., 1993].

In guinea-pig tracheal smooth muscle, SCA40 evoked relaxations that were inhibited by charyldotoxin, suggesting the involvement of BK_{Ca} channels [Laurent et al., 1993]. However, SCA40 had no effect on the activity of BK_{Ca} channels in bovine tracheal SMC [Macmillan et al., 1995]. Cook et al. (1995) reported that relaxant effects of SCA40 in guinea-pig tracheal smooth muscle are due to the inhibition of phosphodiesterase. Activation of K_{Ca} channels in respiratory smooth muscle following β adrenoceptor stimulation appears to be caused by phosphorylation mediated by a cyclic AMP (cAMP)-dependent protein kinase [Kume

et al., 1989]. Thus, the K_{Ca} potassium channel opener (K_{Ca} CO) profile of SCA40 may be due to an indirect action on the cAMP pathway.

Dihydroso-yasaponin I (DHS-I), derived from the medicinal herb *Desmodium adscendens* used therapeutically in the treatment of dysmenorrhea and asthma [Ampofu, 1977], opens charyldotoxin-sensitive K_{Ca} channels in bovine tracheal smooth muscle membranes [McManus et al., 1993]. Data regarding the pharmacology of DHS-I is limited and, therefore, it is not known if it is the active factor of the herb and, if so, whether K^{+} channels are the primary site of action.

The thiazide diuretic hydrochlorothiazide evokes relaxation in vascular smooth muscle due to the activation of charyldotoxin-sensitive K^{+} channels [Calder et al., 1994]. These effects have been associated with an increased $^{86}Rb^{+}$ efflux [Calder et al., 1994] and a decrease in $[Ca^{2+}]_i$ [Pickers and Hughes, 1995]. An increased $^{86}Rb^{+}$ efflux has also been observed with the thiazide drug cyclothiazide in arterial smooth muscle [Moura and Worcel, 1983]. Whether or not thiazides directly act on a K^{+} channel remains to be established.

2.3. Endogenous Openers

Endogenous openers of potassium channels can be divided into three basic groups: (1) ligands that modulate the gating of potassium channels following interaction with plasma-membrane receptors, (2) intracellular second messengers (e.g., inositol triphosphate, inositol 1,3,4,5 tetrakisphosphate, cAMP), and (3) extracellular substances believed to act directly on the channel. Although it is this latter group that will be the subject of this section, the former groups are of great interest in the understanding of the pathophysiology of disease states and subsequent treatment. In pathologic situations, there may be a defect in the neurotransmitter or receptor (e.g., norepinephrine, 5-hydroxytryptamine [5-HT], acetylcholine, opioid peptides), resulting in decreased activation of the associated K^{+} channel. Direct modulation of the K^{+} channel through specific agents could provide focused therapy, depending on the distribution of that channel subtype.

Endothelial cells, which line all blood vessels and the cardiac cavity, have a central role in cardiovascular homeostasis, at least in part, through the release of substances that mediate control of vascular tone and cardiac contraction [Shah, 1992]. Stimuli of receptors (e.g., acetylcholine, bradykinin) or nonreceptor processes (e.g., electrical field stimulation, calcium ionophore A23187) induce endothelium-dependent hyperpolarization of SMC, leading to vasorelaxation [Orlowski et al., 1995].

The first endothelium-derived mediator found to act on vascular SMC was prostacyclin (PGI₂), a vasorelaxing cyclooxygenase product [Weisberg et al., 1978]. The stable analog iloprost, like prostacyclin, evokes a hyperpolarization of SMC through the activation of K^{+} channels [Singel et al., 1990]. K_{Ca} channels have been proposed to be the site of action

of floprost (Boeynaems and Ramboux, 1990); whether this property of floprost is due to activation of PGI₂ receptors or a direct action on the K⁺ channel is still a subject of debate. The development of other stable analogs of PGI₂ that demonstrate therapeutic benefit (e.g., in peripheral arterial occlusive disease, Raynaud's syndrome, cardioprotection, stroke) by the activation of K⁺ channels is an active area of research.

Furchgott and Zawadzki (1980) demonstrated the importance of the vascular endothelium in mediating the relaxant effect of acetylcholine, leading to the proposal of the existence of endothelium-derived relaxing factor (EDRF). EDRF is believed to be nitric oxide (NO) or a closely related NO-releasing molecule (Palmer *et al.*, 1987). EDRF-induced smooth muscle relaxation is mediated, at least in part, by the haem-dependent activation of guanylate cyclase, with the subsequent generation of cyclic GMP (cGMP) (Furchgott and Jothanandan, 1983; Rupprecht and Murad, 1983). Before the discovery of the role of the endothelium in vasodilation, Kuriyama and Suzuki (1978) had observed that acetylcholine hyperpolarized SMC by increasing the membrane permeability to K⁺ ions. Although NO has been reported to change smooth muscle membrane potential (Lanc *et al.*, 1990) and directly activate K_{Ca} in SMC (Bolotina *et al.*, 1994), the contribution of these effects to endothelium-dependent relaxation relative to direct stimulation of guanylate cyclase is unclear. Taylor and Weston (1988) subsequently suggested that an additional factor to EDRF, which could cause vasorelaxation by increasing the membrane potential of the muscle cells, was released from the endothelium by acetylcholine. To distinguish this factor from EDRF, it was termed endothelium-derived hyperpolarizing factor (EDHF). Although evidence has been presented to support the hypothesis that EDHF is a diffusible factor released from the endothelium, an electrotonic spread of hyperpolarization between the endothelial and SMC cannot be eliminated at this stage (Garland *et al.*, 1995). The chemical identity of EDHF remains unresolved; however, a number of candidates (e.g., prostanoids) have been eliminated (Garland *et al.*, 1995). The proposal that EDHF may be a cyclooxygenase P450-derived arachidonic acid metabolite, such as epoxyeicosatrienoic acids, requires confirmation (Gebremedhin *et al.*, 1992; Hecker *et al.*, 1994). Experiments to advance the identification of EDHF may be complicated by the existence of a family of factors, as opposed to a single substance.

The effects of EDHF are associated with the efflux of K⁺ ions from SMC (Chen and Suzuki, 1989), but the channel(s) involved have not been definitively characterized. The majority of studies have found that glibenclamide blocks the change in membrane potential, suggesting the involvement of K_{ATP} channels (Chen and Cheung, 1992; Eckman *et al.*, 1992; Garbino and McPherson, 1992; Adageho and Triggle, 1993). In certain vascular beds, however, K_{Ca} channels have also been proposed to be the site of action of EDHF (Adageho and Triggle, 1993). The existence of more than one EDHF may also complicate the identification of the site of action.

Physiologically, the action of a hyperpolarizing factor that

is distinct from NO appears to predominate in the modulation of endothelial-dependent smooth muscle relaxation in small resistance arteries (Garland *et al.*, 1995). In large arteries, both NO and EDHF appear to contribute to relaxation, with NO being dominant under normal circumstances. Thus, EDHF and the K⁺ channel it activates may be of primary importance in the regulation of vascular resistance. The clinical importance of EDHF in disease aetiology will only be appreciated when the chemical identity is established and/or the K⁺ channel(s) activated is characterized.

The endothelins are a family of 21-amino acid vasoactive peptides, of which endothelin-1 (ET-1) has been reported as the most potent vasoconstrictor known (Randall, 1991). ET-1, however, has also been demonstrated to modulate K_{ATP} channels in both *in vitro* cell culture (Inoue *et al.*, 1990) and *in vivo* studies (Hasegawa *et al.*, 1990; Lippman *et al.*, 1991). As a consequence of this property, ET-1 preferentially evokes vasodilation in certain vascular beds (e.g., pulmonary; Lippman *et al.*, 1991). In addition, the vascular effects of benzopyran, but not carbocyclic and pyridyl-quinazolinone, K_{ATP}CCs (see Section 2.1) were modified by ET-1, suggesting that BRL-38227 and ET-1 have affinity for a common site on the K_{ATP} channel (Lawson *et al.*, 1992). This hypothesis is supported by the finding that BRL-38227 inhibits binding of [¹²⁵I]ET-1 to rat cardiac membranes (Waugh *et al.*, 1992). Further, ET-1 evokes membrane hyperpolarization in rat aortic SMC (Van Renterghem *et al.*, 1988), rat glioma C6-BU-1 cells (Cleaveson *et al.*, 1991), porcine coronary artery cells (Hu *et al.*, 1991), rat gastric longitudinal and circular smooth muscle (Fulginiti *et al.*, 1993) and guinea-pig tracheal smooth muscle (Lévesque *et al.*, 1991) through the activation of charybdotoxin-sensitive K_{Ca} channels. In contrast, ET-1 also exhibits antagonistic properties at both K_{ATP} (Miyoshi *et al.*, 1992) and BK_{Ca} (Hu *et al.*, 1991) channels.

Consequently, ET-1 can directly modulate K⁺ channels having affinity for both K_{ATP} and K_{Ca} channels. The (patho)physiological role of ET-1-induced K⁺ channel activation remains to be established. The recent advances in the identification of selective antagonists will allow the elucidation of the involvement, if any, of ET-receptors in this action of the peptide.

Interestingly, ET-1 and ET-2 also evoke an indirect endothelium-dependent hyperpolarization in rat mesenteric smooth muscle (Nakashima and Vanhoutte, 1993a), but not in canine coronary artery (Nakashima and Vanhoutte, 1993b).

Finally, a variety of endogenous polypeptides have been proposed to exhibit K⁺ channel opening properties. The endogenous vasodilator peptides, vasoactive intestinal peptide and calcitonin gene-related peptide (CGRP), activate K_{ATP} channels in vascular SMC, leading to hyperpolarization and tissue relaxation (Standen *et al.*, 1988; Nelson *et al.*, 1990). CGRP is a polypeptide located in neurones that form a close association with both central and peripheral blood vessels (Bevan and Brannen, 1986). The CGRP-induced relaxations of some blood vessels (e.g., rat aorta) are endothelium-dependent; thus, the role of endothelium-

derived factors in the observed hyperpolarization has been proposed (Grace et al., 1987; Nelson et al., 1990). Galanin and somatostatin, hyperglycemic hormones, activate K_{ATP} channels in insulin-secreting pancreatic β -cells (De Wille et al., 1988, 1989). The block of anoxia-induced depolarization of hippocampal CA3 neurones and glutamate release inhibition by galanin is consistent with K^+ channel opening (Ben-Ari, 1990). However, not all reports support the conclusion of K^+ channel activation as a property of vasoactive intestinal peptide, CGRP, galanin, and somatostatin; further work is required to establish the role of K^+ channels in their physiological profiles.

2.4. Others

1,1-Dimethyl-4-phenylpiperazinium (DMPP), a selective nicotinic receptor stimulant, has shown activity consistent with K^+ channel activation in the isolated tunica muscularis mucosae of the rat oesophagus (Adenambo et al., 1993). Neither apamin nor glyburide modified the DMPP-induced relaxation. Whether this was a direct action of DMPP on a K^+ channel or a nicotinic-choolinergic-linked K^+ channel requires elucidation.

3. MECHANISM(S) OF ACTION

The standard criteria that initially identified a compound as a KCOs has been its ability to relax an *in vitro* smooth muscle preparation contracted with low, but not high, concentrations of extracellular K^+ ions (Weir and Weston, 1986; Bray et al., 1987; Hamilton and Weston, 1989; Lawson and Cavero, 1989). The principal determinant for inhibition or reversal of smooth muscle contraction by a KCO is a reduction in the free cytosolic Ca^{2+} concentration. The mechanism(s) involved in the production of smooth muscle relaxation by K^+ channel openers, especially K_{ATP} COs, has been (and still is) a major subject of debate. To date, information regarding mechanism(s) of action primarily has been obtained from studying K_{ATP} COs, especially cromakalim and other benzopyrans. Investigations with other classes of KCOs (e.g., K_v COs) that will demonstrate whether or not mechanism(s) of action are common between the different K^+ channels and how they are activated by different ligands, are awaited. The mechanism of action involved in a given response may not only depend on the subtype of K^+ channel, but also how the KCO interacts with the channel to evoke that effect (Lawson et al., 1992; Randall and Griffith, 1993).

Although some mechanism(s) are universal in different excitable and nonexcitable cells, not all processes will be applicable to all types of tissue. The majority of data has been obtained from SMC and requires confirmation in other cells.

3.1. Hyperpolarization

The opening of K^+ channels by these compounds and subsequent efflux of K^+ ions from the cytosol leads to mem-

brane repolarization and/or hyperpolarization (Cook, 1990; Edwards and Weston, 1990; Robertson and Steinberg, 1990; Longman and Hamilton, 1992). This change in membrane potential is followed by a reduction in cytosolic free Ca^{2+} and/or an inhibition of mechanisms producing increases in cytosolic free Ca^{2+} . The outcome of these effects is a reduction in membrane and cell excitability, resulting in a greater cellular resistance to activation by excitatory stimuli.

In smooth muscle preparations, the relaxant actions of K_{ATP} COs have been accompanied by an increase in negativity of the resting membrane potential (hyperpolarization) towards the calculated E_K , together with an outward current of K^+ ions (Hamilton and Weston, 1989).

An excellent correlation exists between the potencies of the K_{ATP} COs for stimulation of $^{86}Rb^+$ efflux and vasorelaxation in the aorta and portal vein *in vitro* preparations (Quast et al., 1992), supporting the hypothesis that the functional responses rely on the opening of plasmalemma K^+ channels in vascular smooth muscle.

Although, initially, it was assumed that the hyperpolarization caused by K^+ efflux produced closure of VOCs, preventing depolarization-induced Ca^{2+} entry into the cell, other mechanisms, in light of recent evidence, may also contribute to the effects produced by K^+ channel openers (see Sections 3.2, 3.3 and 3.4). Evidence that cromakalim-evoked increase in $^{86}Rb^+$ efflux or hyperpolarization of vascular smooth muscle tissue was not influenced by lanthanum or the Ca^{2+} antagonist nifedipine, is indicative that the action of the K_{ATP} COs is not dependent upon modifying the influx of external Ca^{2+} ions (Coldwell and Howlett, 1988; Southerton et al., 1988).

Vasorelaxant effects of K_{ATP} COs have been reported that are independent of membrane hyperpolarization or ion efflux (Hamilton et al., 1986; Quast and Baumbach, 1988; Greenwood and Weston, 1991). Such findings suggest that these drugs can exert a response through mechanisms other than the opening of K^+ channels. The possibility that K_{ATP} COs do not interact directly with an ion channel but, rather, for example, with an enzyme system involved in intracellular phosphorylation, provides a novel explanation for some of the apparently anomalous effects of these agents (Edwards and Weston, 1994).

3.2. Intracellular Ca^{2+} Stores

Experimental evidence suggests that K_{ATP} COs may also have a direct effect on intracellular stores. Cromakalim evoked a constrictor response in rabbit aorta bathed in a Ca^{2+} -free solution, which may be related to effects on intracellular Ca^{2+} stores (Duty and Weston, 1992). Using rabbit cultured tracheal SMC, Chopra et al. (1992) demonstrated that cromakalim reduced the uptake into and inhibited the release of $^{45}Ca^{2+}$ from the sarcoplasmic reticulum. These findings support those obtained from vascular smooth muscle, where contractile responses to noradrenaline, dependent on intracellular calcium stores, were attenuated by cromakalim (Bray et al., 1991).

In contrast, the effect of cromakalim on rat pulmonary artery did not appear to involve an action on Ca^{2+} release from internal stores (Savineau and Marthan, 1993).

3.3. Inositol Phosphate Cascade

BRL 18227 inhibited Ca^{2+} release from intracellular stores of rabbit isolated mesenteric artery due to an action on noradrenergically-stimulated 1,4,5-inositol triphosphate production (Ho et al., 1991). This effect was sensitive to glibenclamide and high (128 mM) extracellular KCl. Hyperpolarization of the plasma membrane of canine coronary artery by K_{ATP} COs has also been associated with an inhibition of the production of 1,4,5-inositol triphosphate and, hence, Ca^{2+} release from intracellular stores (Yamagishi et al., 1992). Interestingly, K_{ATP} COs have shown no effect upon phosphatidylinositol turnover in brain tissue (Coldwell and Howlett, 1988).

3.4. Ca^{2+} Sensitivity

The hyperpolarization by K_{ATP} COs (e.g., levocromakalim) may also be linked with a reduction in the sensitivity of the contractile elements of vascular smooth muscle to Ca^{2+} (Okada et al., 1993). Cromakalim, however, has no direct effect on the contractile proteins of skinned rat aortic muscle (Allen et al., 1986).

3.5. Others

Cromakalim has been reported to stimulate the Na-K pump in human and canine mesenteric arteries through an elevation in intracellular Na⁺ (Giong et al., 1993). In rat aorta, cromakalim appears to be absent of effects on the Na-K pump (Cavero et al., 1987).

The K_{ATP} COs, cromakalim, and RP-49356 failed to increase cAMP or cGMP in SMC (Southerton et al., 1988; Nakajima et al., 1989), or to potentiate the effects of forskolin in airway smooth muscle preparations (Berry et al., 1991; Murray et al., 1990). Pincicidil and cromakalim do not appear to be inhibitors of phosphodiesterase activity in airway smooth muscle (Berry et al., 1991; Ho et al., 1990). Guanine nucleotide-binding proteins (G-proteins) are known to provide a link between membrane receptors and intracellular events, and the possibility exists that G-proteins may modulate the activity of a K^{+} channel sensitive to K_{ATP} COs. Results with pertussis or cholera toxin, however, suggest that the corresponding G-proteins are not involved in the actions of cromakalim in vascular smooth muscle (Longman and Hamilton, 1992).

Therefore, the involvement of intracellular Ca^{2+} stores in the responses to K_{ATP} COs may be species- and/or tissue-dependent. Further studies are required to determine whether or not this is a property unique to cromakalim (or benzopyrans) and if it is a direct effect or a consequence of hyperpolarization. Thus, the mode of action of K^{+} channel openers may not be as simple as first thought and more research effort is required in this area.

4. THERAPEUTIC TARGETS AND POTENTIAL

Theoretically, at least, the general decrease in the excitability of cells that follows K^{+} channel opening infers a broad clinical potential for drugs with this property in a number of pathologic conditions.

4.1. Cardiovascular System

The preclinical profile of K_{ATP} COs clearly supports a clinical potential for their use in vascular pathologies that require a decrease in peripheral vascular resistance, an inhibition of excessive vasoconstriction, and/or a prolongation of myocardial tissue viability, while undergoing transient oxygen deficiency.

4.1.1. Vascular. The pharmacological profiles of K_{ATP} COs in vascular smooth muscle tissues and in vivo models of vascular disorders (e.g., hypertension, peripheral vascular disease, angina) have been reviewed extensively (Edwards and Weston, 1990; Richer et al., 1990; Longman and Hamilton, 1992; Edwards and Weston, 1995). In a variety of vascular tissues from a range of species, K_{ATP} COs display the ability to relax smooth muscle by inhibiting both spontaneous tone and/or spasmogen-induced contraction. Arterial smooth muscle tone, of which K^{+} channels in the SMC are important regulators, is the main determinant of peripheral vascular resistance and blood pressure. Functional defects of K^{+} channels may lead to vasoconstriction or compromise the ability of an artery to dilate in pathologic conditions of the vasculature, such as vasospasm, hypertension, ischemia, hypotension following sepsis, and diabetes.

Activation of K_{ATP} channels, which respond to the metabolic state of the cell, may be involved in arterial dilation in reactive hyperemia, septic shock, ischemia, and hypoxia (Nelson, 1993). The involvement of K_{ATP} channels in these clinical conditions may not be exclusive; however, the absence of selective agents of other K^{+} channels presently limits full characterization of the aetiology of such diseases.

K_{ATP} COs can lower both normal and experimentally elevated blood pressure (Edwards and Weston, 1990; Richer et al., 1990; Longman and Hamilton, 1992). The mechanism of this effect is an actively mediated decrease in total peripheral vascular resistance. Therapeutic experience with pincicidil, diazoxide, and minoxidil indicates that they also reduce elevated blood pressure in patients (Gross, 1977; Oates et al., 1977; Ahlberg-Remue, 1985). Hypertensive patients treated with pincicidil (10–25 mg b.i.d.) showed a marked fall in total peripheral resistance, the short duration of which was controlled through the development of sustained-release formulations (Carlson et al., 1983, 1985). The antihypertensive effects of pincicidil are accompanied by an initial reflex tachycardia and by weight gain in approximately 40% of patients. Although the antihypertensive effects of diazoxide have been known for many years (Gross, 1977; Oates et al., 1977), increased blood glucose levels has restricted its use to hypertensive emergencies. The extent to which the antihypertensive effects of diazoxide result from K^{+} channel

opening action is still unknown, as evidence of additional effects have been observed (Newgreen et al., 1990). The relatively high incidence of fluid retention has severely restricted the clinical use of nitroglycerin (Gross, 1977; Oates et al., 1977). The K^+ channel opening profile of nitroglycerin, however, is different from that of diazoxide and cromakalim, whereby the involvement of a nonselective channel has been proposed (Newgreen et al., 1990; Lawson and Hicks, 1993).

Clinical experience with cromakalim is less extensive than for the above K_{ATP} COs, and there are no published reports of clinical trials with the other K_{ATP} COs (e.g., RP 49356, Ro 31-6930, EMD 52692, KRN 2391). In mild to moderate hypertension, cromakalim (0.5–1.5 mg po) lowered systolic and diastolic blood pressure following a single oral dose (Vanden Burg et al., 1986; Singer et al., 1989) or chronic once-daily administration (Eckel and Greb, 1987; Lebel et al., 1989; Vanden Burg et al., 1987), with no such effects in a parallel normotensive group or when placebo was administered. As a direct vasodilator, the use of KCOs as monotherapy to reduce blood pressure can produce a series of undesirable effects (e.g., tachycardia, headache, flushing, increase in renin, aldosterone, and catecholamine secretion, and sodium and water retention) (Glück et al., 1987; Lijnen et al., 1989) that are not acceptable in clinical practice. K_{ATP} COs, however, could become useful antihypertensive therapy if appropriately formulated and co-prescribed with selected agents to reduce or prevent the undesirable events. Due to apparent cardioprotective properties (see Section 4.1.2), small doses of K_{ATP} COs could be used to provide myocardial protection to hypertensive patients, an area where current drugs do not appear to substantially reduce cardiovascular mortality and morbidity (Escande and Caverio, 1992; Caverio and Premiersart, 1994; Grover, 1994a,b).

In preclinical studies, K_{ATP} COs relaxed coronary conductance arteries, increased selectively coronary blood flow and antagonized the vasoconstrictor activity of a large number of excitatory stimuli (Longman and Hamilton, 1992). If similar findings are reproduced in human subjects, K_{ATP} COs would have antianginal activity at doses that do not provoke undesirable, reflex-mediated activation of the sympathetic system, which is a deleterious physiological reaction for myocardium frankly ischemic or lacking a safety margin of blood flow reserve. Thus, these agents demonstrate properties desirable to improve oxygen delivery and also reduce oxygen consumption within ischemic regions of patients with transient and chronic heart disease (e.g., angina pectoris). Reports on the effects of KCOs in angina pectoris, however, are restricted primarily to trials in which nicorandil was used (Fukushima, 1993; Kuroshita and Sakai, 1990). The clinical benefits of nicorandil probably result both from K^+ channel opening properties and its ability to stimulate smooth muscle guanylate cyclase (Hamilton and Weston, 1989). Cromakalim may also be beneficial in the treatment of angina pectoris (Thomas et al., 1990).

One therapeutic approach to congestive heart failure, although it is of a symptomatic nature, is to reduce peripheral vascular resistance, a property demonstrated by

K_{ATP} COs (Gopalakrishnan and Triggle, 1990). In an *in vivo* rat cardiac failure model, down-regulation of ventricular K_{ATP} channel density, together with 1,4-dihydropyridine-sensitive Ca^{2+} channels and β -adrenoceptor densities, was observed, and these changes have been implicated in this pathophysiological condition (Gopalakrishnan et al., 1990). No clinical data for the effectiveness of KCOs in the treatment of congestive heart failure has been published.

Activation of K^+ channels can improve the energy metabolism and the mechanical performance of skeletal muscles suffering oxygen deficiency (Cook and Chapman, 1993). This is achieved partly by a selective dilation of collateral vessels supplying the ischemic skeletal muscle and, partly, by a better utilization of high energy phosphates. In rat skeletal muscle, Angersbach and Nicholson (1988) demonstrated that K_{ATP} COs, but not Ca^{2+} antagonists or hyalazine, selectively increased blood flow to collateral vessels in a previously ischemic limb, despite a reduction in basal diastolic blood pressure. These mechanisms are evidently of therapeutic potential for treating patients with peripheral vascular disease, a disabling old-age disorder characterized by poor blood supply to the limbs due mostly to atherosclerosis. The role of K^+ channel activation in the beneficial effects of isoproterenol (see Section 2.3; Pessi et al., 1986; Müller Buhl et al., 1987; Oberender et al., 1989) in peripheral arterial occlusive disease still remains to be determined.

Cromakalim (10 μ M) increased $^{86}Rb^+$ efflux in control cultured arterial SMC, but was without effect in cholesterol-enriched SMC (Ihlenko and Bialecki, 1989). Cholesterol enrichment of SMC membranes can severely influence the cellular responses to cromakalim. Thus, the benefit of K_{ATP} COs in clinical conditions associated with excess lipids may be questionable.

K_{Ca} channels appear to play a fundamental role in regulating the degree of intrinsic tone in resistance arteries (Nelson, 1993) and, as such, help regulate arterial responses to pressure and vasoconstrictors. Therefore, activation of these channels should contribute to vasodilation. Defects in K_{Ca} channels could lead to, or contribute to, pathologic conditions that are characterized by highly constricted arteries. Thus, the development of activators may be useful in the treatment of, for example, coronary or cerebral vasospasm. Although NS-004 and NS-1619 activate HK_{Ca} channels (see Section 2.2), their relaxant effects in rat portal vein (Edwards et al., 1994) and cardioprotective effects (Sargent et al., 1993) are probably due to inhibition of L-type calcium channels.

As blood flow increases through a conduit artery, the vessel dilates (Hilton, 1959) by an endothelium-dependent mechanism (Hail et al., 1986; Fohl et al., 1988). Flow in rabbit isolated thoracic arteries appears to activate a charyldotoxin-sensitive K^+ channel on the endothelial cell membrane that leads to the release of NO (Cooke et al., 1993). Neither glibenclamide (K_{ATP} channel blocker) nor apamin (SK_{Ca} channel blocker) had any effect on the flow-mediated vasodilation. Thus, in the regulation of arterial tone, HK_{Ca} channels act as the transducer of the flow stimulus, whereas NO is the effector of the vasodilation.

Endothelium-dependent vasodilations have also been associated with the activation of K_{ATP} channels. The cromakalim- and pinacidil-induced dilation of canine large coronary arteries *in vivo*, an indirect flow-mediated effect, are entirely dependent on the endothelium (Drieu La Rochelle *et al.*, 1992; Ghaleb *et al.*, 1993). The $K_{ATP}CO$, levcromakalim, was more potent as a vasorelaxant in rat aortic ring preparation with endothelium than in denuded tissues, an effect that involved NO (Lawson *et al.*, 1993b).

In porcine aortic endothelial cells, pinacidil and cromakalim elevated $[Ca^{2+}]_i$ by inducing membrane hyperpolarization following K^+ channel activation (Lackhoff and Busse, 1990), an effect that can promote Ca^{2+} -dependent formation of EDHF. Pinacidil also opened K_{ATP} channels in both rat aorta and brain microvascular endothelial cells (Jang *et al.*, 1993). These findings suggest that K_{ATP} channels may play a role in the regulation of endothelial cell resting membrane potential, for example, during impaired energy supply and, therefore, modulate release of endothelium-derived vasoactive factors and blood flow.

4.1.2. Cardiac. K^+ channel opening properties are desirable for therapeutic agents aimed at treating patients with transient and chronic heart diseases. $K_{ATP}CO$ s have demonstrated cardioprotective effects as a consequence of improving oxygen delivery and reducing oxygen consumption within the ischemic region (Escande and Caverio, 1992; Grover, 1994a,b; Yao and Gross, 1994; Grover *et al.*, 1995). Both antiarrhythmic and proarrhythmic properties have been reported in $K_{ATP}CO$ s, leading to the safety of these drugs being a major subject of discussion (Carlsson *et al.*, 1991; Tosaki *et al.*, 1993; Black and Lucchesia, 1994; D'Alonzo and Grinvald, 1994; Wilde, 1994). Although there appear to be good theoretical arguments as to why $K_{ATP}CO$ s may be of use in the treatment of some arrhythmias and in ischemic heart disease, there are major hurdles to overcome.

Clinical evidence to establish the benefits of $K_{ATP}CO$ s as treatments of patients with heart disease is awaited. Therefore, only the basic concepts of $K_{ATP}CO$ -induced beneficial or undesirable cardiac effects will be outlined.

Cardiac K_{ATP} channels have been shown to open in response to ischemia (Kantor *et al.*, 1990). The depletion of ATP in the myocardium and subsequent opening of K_{ATP} channels may lead to a rapid reduction in contractility of the ischemic myocardium to protect against further ischemic injury. In support of this hypothesis, cromakalim, RP 52891 (aprikalim), and pinacidil, in animal models, cause a glibenclamide-sensitive reduction in the severity of ischemic/reperfusion injury of the myocardium (Bicher *et al.*, 1990; Auchincloss *et al.*, 1992; Escande and Caverio, 1992; Grover, 1994a,b). Thus, $K_{ATP}CO$ s play a cardioprotective role, whereas glibenclamide worsens myocardial stunning (Auchincloss *et al.*, 1992). Analogies have been observed between the cardioprotection conferred by $K_{ATP}CO$ s and ischemic preconditioning (*i.e.*, increased tolerance of cardiac myocytes to an ordinarily lethal ischemic insult, achieved by an initial brief exposure to ischemia), for example, both

are sensitive to glibenclamide blockade (Gross and Auchincloss, 1992). Thus, therapy with $K_{ATP}CO$ s may afford a permanent 'chemical preconditioning' that confers on the heart the ability to better withstand transient oxygen deprivation and, consequently, to suffer less tissue damage during acute myocardial infarction.

Whether $K_{ATP}CO$ s exert their beneficial effects on the ischemic heart by a direct (myocardial) or indirect (vascular) action remains to be determined. Studies with U-89,232 (cromakalim analog; Foombs *et al.*, 1992) and BMS 180446 (pinacidil analog; Grover *et al.*, 1995), $K_{ATP}CO$ s devoid of vascular effects demonstrated cardioprotection against ischemia in animal models greater than that observed with cromakalim. This would suggest that there is a direct myocardial action of these second generation $K_{ATP}CO$ s that will provide an opportunity to explore the cardioprotection of such agents without possible complication (*e.g.*, hypotension, coronary steal). The mechanism of tissue selectivity is not clear, but it may be related to the existence of $K_{ATP}CO$ 'receptor' subtypes in different tissues (Attwell, 1994). Similar concepts of subtypes of the site(s) of action of $K_{ATP}CO$ s in SMC previously have been proposed in SMC (Piper *et al.*, 1990; Wickenden *et al.*, 1991; Lawson and Hicks, 1993).

Although K_{ATP} channel opening in response to ischemia may offer a cardioprotective mechanism, there is another consequence. The resulting increase in K^+ efflux shortens the action potential duration and contributes to the extracellular K^+ accumulation observed during an ischemic episode (Coetzee, 1992). These changes in conductance and extracellular K^+ have been hypothesized to be responsible for ischemia-induced arrhythmias. $K_{ATP}CO$ s have demonstrated arrhythmogenic properties in certain animal studies (Chi *et al.*, 1990; Tosaki *et al.*, 1992; De La Coussaye *et al.*, 1993). However, although $K_{ATP}CO$ s may be contraindicated in some types of arrhythmia, they may be of use in the treatment of certain other types of arrhythmia resulting from a repolarization defect (Antzelevitch and Di Diego, 1992). $K_{ATP}CO$ s have been shown to suppress rhythm abnormalities related to delayed repolarization and early afterdepolarizations in anaesthetized rabbits (Carlsson *et al.*, 1991).

4.1.3. Blood. The $K_{ATP}CO$ s, cromakalim, celikalim, and pinacidil, inhibited white thrombus formation in a rabbit arteriovenous shunt model, although they had no effect on human platelet aggregation (Paredas *et al.*, 1994). Antithrombotic activity of $K_{ATP}CO$ s *in vivo* may be related to beneficial effects on blood rheology and reduced red blood cell deformability.

4.2. Respiratory System

Administration of cromakalim and other $K_{ATP}CO$ s to conscious (oral or inhalation route) or anaesthetized (oral, inhalation or *i.v.* route) guinea-pigs protects against histamine, 5-HT, or (in sensitized animals) ovalbumin induced bronchoconstriction (Ruebner and Karlsson, 1993). In the anaesthetized animal (Konzen-Russler model, where protective cardiovascular reflexes are inhibited), a reduction in

diastolic blood pressure was observed following oral or i.v. administration of $K_{ATP}COs$. Bronchodilation, however, could be achieved at doses of cromakalim not reducing mean arterial blood pressure in experiments where the $K_{ATP}CO$ was administered by inhalation; thus, demonstrating selectivity (Bowring *et al.*, 1991; Raeburn and Karlsson, 1991; Bowring *et al.*, 1993; Arch *et al.*, 1994). Respiratory dynamics measurements in anaesthetised guinea-pigs revealed that the $K_{ATP}COs$, cromakalim and BRL 55834, resembled theophylline by eliciting similar inhibition of histamine-induced increase in airways resistance and decrease in lung compliance, and salbutamol, a β -agonist, was more effective against resistance than compliance (Bowring *et al.*, 1990). Therefore, $K_{ATP}COs$, compared to β -agonists are more effective dilators of small airways (where constriction decreases compliance) for identical large airways effects (where constriction increases resistance). Reports have implicated the activation of K_{Ca} channels in the relaxant responses of respiratory smooth muscle to β -agonists (Kume *et al.*, 1989; Jones *et al.*, 1990). Thus, these findings may be suggestive of the distribution of K_{ATP} and K_{Ca} channels within the smooth muscle of the respiratory system. The clinical relevance of such a hypothesis is, as yet, unclear.

$K_{ATP}COs$ can inhibit neurotransmitter release from cholinergic and nonadrenergic, noncholinergic excitatory neurones (NANCs) in guinea-pig lung *in vitro* and *in vivo* (Raeburn and Karlsson, 1991; Small *et al.*, 1992). These neural effects of $K_{ATP}COs$ may be very relevant to potential treatment of asthma because, not only does bronchoconstriction frequently have a significant parasympathetic component, but also neurogenic inflammation of the lung may contribute to the pathology of asthma (Barnes *et al.*, 1991). The main evidence for an effect on neurotransmitter release is that $K_{ATP}COs$ are far more effective at inhibiting cholinergically or NANC-mediated bronchoconstriction or mucus secretion elicited by stimulating neurotransmitter release than when the relevant neurotransmitter is supplied directly (Ichinose and Barnes, 1990; Burka *et al.*, 1991; Raeburn and Karlsson, 1991; Small *et al.*, 1992). A prejunctional site of action has been proposed for the inhibition of peptidergic excitatory neurotransmitter release due to $K_{ATP}COs$ inhibiting the NANC neurotransmitter transmission at concentrations slightly lower than those causing relaxation of airways smooth muscle. Interestingly, the $K_{ATP}COs$ do not seem to interfere with NANC inhibitory neurotransmitter transmission in the lung (Burka *et al.*, 1991). Nielsen-Kudsk *et al.* (1994) demonstrated that, like cromakalim and pinacidil, terbutaline (β -agonist), theophylline (xanthine), and verapamil (Ca^{2+} antagonist) induced inhibition of NANC neurotransmission in guinea-pig bronchi involved a prejunctional site.

The inhibition by cromakalim of electrically evoked [H]-acetylcholine release in rat isolated trachea has been suggested to be an epithelium-dependent mechanism (Weisler *et al.*, 1993). This effect was only observed in tube preparations, where the mucosal/submucosal environment would be better preserved, and not in trachea opened longitudinally.

$K_{ATP}COs$, BRL 38227 and YM-934, inhibited plasma leakage in trachea, main bronchi, and central and peripheral intrapulmonary airways evoked by stimulation of vagal nerves in guinea pigs (Lei *et al.*, 1993; Ishikawa *et al.*, 1994). These compounds had no effect on exogenously administered Substance P-induced plasma leakage. Thus, cromakalim and YM-934 inhibit airway neurogenic inflammation by modulating the release of neuropeptides from the sensory nerve endings, and the inhibitory effect can be attributed to the KCO activity.

$K_{ATP}COs$ can reduce obstruction to airflow by suppressing hyperactivity of intact airways in animals, with doses that are insufficient to relax airway smooth muscle *in situ* in normal animals (Chapman *et al.*, 1991; Paciorek *et al.*, 1992; Morley, 1994). Hence, the potency of $K_{ATP}COs$ as inhibitors of bronchoconstriction is greater in hyperreactive than normal animals. An almost universal characteristic of asthmatics is that their airways are hyperresponsive to a wide range of physiological and pharmacological stimuli (Smith, 1992). The causes of airways hyperresponsiveness in humans are not well-defined, although several animal models have been developed to emulate this response. In general, however, the degree of hyperresponsiveness achieved in animals is less than that in humans (Smith, 1989). Compounds that open K^+ channels and impair expression of airway hyperactivity in the absence of direct smooth muscle spasmolysis will provide a novel approach to symptomatic therapy in asthma (Morley, 1994).

In general, the direct relaxant properties of $K_{ATP}COs$ such as cromakalim have been assessed predominantly in guinea-pig isolated trachealis muscle (Raeburn and Karlsson, 1991; Longman and Hamilton, 1992; Small *et al.*, 1992). This tissue has also been used for ion flux studies and for intracellular recording of change in membrane potential. $K_{ATP}COs$ inhibit contractions to or reverse precontractions to a variety of spasmogens in guinea-pig tracheal preparations. However, in interaction studies, the smooth muscle relaxant responses to $K_{ATP}COs$ in the guinea-pig trachea are blocked by glibenclamide in a manner that is not consistent with competitive antagonism (*i.e.*, the maximum effect of the $K_{ATP}CO$ is reduced in the presence of the antagonist [Berry *et al.*, 1991; Nielsen-Kudsk *et al.*, 1993]), but is consistent with a lack of spare channels (*i.e.*, the $K_{ATP}CO$ must activate all channels in the tissue to evoke a maximal response; see Section 2.1). The antagonism by glibenclamide of the effects of $K_{ATP}COs$ in functional *in vitro* pharmacology studies on vascular smooth muscle and certain nonvascular smooth muscle preparations has been described as competitive in nature (see Section 2.1).

The lack of competitive interaction between glibenclamide and the $K_{ATP}COs$ is indicative of the involvement of more than one mechanism in the relaxant effect of the latter. This is consistent with the conclusion that relaxations of tracheal smooth muscle to BRL 55834 (a benzopyran $K_{ATP}CO$) is mediated by, at least, a glibenclamide-sensitive and a glibenclamide-resistant K^+ channel (Lawson *et al.*, 1993a; Edwards *et al.*, 1995). In addition, BRL 55834 has

been reported to activate an ATP- and glibenclamide-sensitive K^+ channel and, at higher concentrations, a large conductance charybdotoxin-sensitive Ca^{2+} -activated K^+ channel in bovine trachealis SMC (Ward *et al.*, 1992).

Cromakalim and levocromakalim have demonstrated relaxant effects in human bronchial smooth muscle (Taylor *et al.*, 1992; Black *et al.*, 1990). Differences (primarily in potency) from the findings obtained in guinea-pig preparations suggested that the guinea pig is not a good predictor of the inhibitory response of $K_{ATP}COs$ in humans.

Clinical trial of cromakalim in patients with nocturnal asthma showed that a single (0.5 mg) or repeat (0.25, 0.5 mg) oral dose administered at 11:00 p.m. could attenuate the dip in lung function measured at 6:00 a.m. the following morning (Williams *et al.*, 1990). The predicted peak plasma concentration of cromakalim in these studies was about 5-fold less than its threshold concentration for relaxation of tone in human bronchi (Taylor *et al.*, 1992). Studies in animal models prior and subsequent to these findings suggest that the positive results are due to actions other than just relaxation of the bronchial smooth muscle (Longman and Hamilton, 1992; Small *et al.*, 1992; Morley, 1994). The efficacy of cromakalim may not involve the direct relaxation of airway smooth muscle, but is due to an influence on neural mechanisms underlying airway hyperresponsiveness. This suggestion is supported by the effects of $K_{ATP}COs$ in animal models of hyperreactivity (Morley, 1994); the potency of cromakalim in human hyperresponsive airway smooth muscle is, as yet, unknown.

Interestingly, when the dose of cromakalim was increased to 1.5 mg (single dose), no significant reduction in the morning dip in lung function was observed (Williams *et al.*, 1990). The failure of the latter dose of cromakalim to improve lung function was attributed to 10 out of the 23 subjects being unable to exert maximal expiratory effort during measurements of FEV₁ values (forced expiratory volume in 1 sec) as a consequence of headache. BRL 38227, which replaced cromakalim in clinical trials, failed to elicit significant bronchodilation or reduce bronchial hyperresponsiveness to histamine or methacholine when administered as a single oral dose to asthmatics (Kidney *et al.*, 1993); thus, not meeting the criteria set for its development in asthma. As with cromakalim, the dose-limiting side effect of oral BRL 38227 is headache, probably resulting from cerebral vasodilation (Arch *et al.*, 1992). Bimatolol, an analog of cromakalim, also lacked bronchodilatory effects following inhaled administration to mild to moderate bronchial asthmatic adult patients (Rauchoy *et al.*, 1994). Whether this was a true lack of bronchial efficacy or that the dose of drug, to avoid other effects (no headache or cardiovascular effects reported), was too low requires further investigation. Therefore, to be useful oral bronchodilators, and have the potential to reduce bronchial hyperresponsiveness, $K_{ATP}COs$ with selectivity for airway relative to vascular smooth muscle greater than that of cromakalim or BRL 38227 are required.

BK_{Ca} have been demonstrated in high density in canine, bovine, and human airway smooth muscle (McCaig and

Welsh, 1986; Green *et al.*, 1991; Miura *et al.*, 1991). This has led to the proposal that openers of BK_{Ca} channels could demonstrate therapeutic benefit in regulation of tone of the respiratory system. The findings involving BK_{Ca} channels in the effects of β -agonists on airway smooth muscle (Kumar *et al.*, 1989; Jones *et al.*, 1990) support this hypothesis. In human bronchi, however, NS 1619 (a BK_{Ca} channel opener) (Olesen *et al.*, 1994a,b) evoked a weak relaxant effect that did not involve BK_{Ca} channels, which could call into question the utility of such drugs as bronchodilators (Templeton *et al.*, 1993). The lack of selectivity of $K_{Ca}COs$ for the BK_{Ca} channel (see Section 2.2) has delayed progress in this area and, thus, appreciation of the true utility of such agents.

4.3. Reproductive System

By virtue of the smooth muscle relaxing effects, K_{ATP} channel openers may be useful in the treatment of premature labour and dystocicorrhoea (Piper *et al.*, 1990). Several $K_{ATP}COs$ are capable of producing glibenclamide-sensitive relaxation of uterine smooth muscle of rat, both *in vitro* and *in vivo* (Piper *et al.*, 1992). Cromakalim inhibited the spontaneous phasic activity and spasmodic-induced contractions of isolated uterus from the term-pregnant rat (Hollingsworth *et al.*, 1987). BRL 38227 and pinacidil inhibited spontaneous and oxytocin-induced contractions in human isolated pregnant myometrium, obtained before and after the onset of labour (Cheuk *et al.*, 1993; Morrison *et al.*, 1993). The $K_{ATP}COs$ were more potent in nonpregnant than pregnant human myometrium (Cheuk *et al.*, 1993). The relaxant effects of the two $K_{ATP}COs$ in human pregnant myometrium was sensitive to glibenclamide. Thus, $K_{ATP}COs$ may have potential as a new generation of tocolytic agents. Preferential higher potency would suggest that $K_{ATP}COs$ would be more effective tocolytic agents in nonpregnant than pregnant women (Cheuk *et al.*, 1993). Not all women with preterm uterine contraction, however, are candidates for tocolysis (Monga and Casas, 1995).

Although channels permeable to Rb^+ and K^+ exist in the uterus, cromakalim does not stimulate efflux of these ions in rat uterus, even though relaxant responses were sensitive to glibenclamide (Hollingsworth *et al.*, 1987). This finding was further supported in humans isolated myometrium, where Rb^+ exhibited differentiation on the effects of BRL 38227 and P1060 to pinacidil analog on amplitude and frequency of spontaneous contraction (Criddle and Soares de Moura, 1995). In these studies, Rb^+ -sensitive and Rb^+ -insensitive mechanisms were identified, of which the former appeared more important in efforts of P1060 than BRL 38227. Although $K_{ATP}COs$ may demonstrate benefit in uterine-associated disorders, the involvement of K^+ channels still requires confirmation.

β -Adrenoceptor agonists, relaxin, and other uterine relaxants that increase intracellular cAMP levels, activate K_{Ca} channels in myometrial cells (Trentham *et al.*, 1991; Meera *et al.*, 1994; Saborin, 1995). Ibexotoxin, a K_{Ca} channel blocker, depolarizes myometrial cells and increases phasic

contractions in rat and human myometrial preparations (Anwer *et al.*, 1993). Therefore, selective direct openers of K_{Ca} channel (iberiotoxin-sensitive) may also have therapeutic benefit in uterine-associated disorders.

Interestingly, DHS-3, derived from the medicinal herb *Desmodium adscendens* and used therapeutically in the treatment of dysmenorrhea (Anpofo, 1977), opens charybdotoxin-sensitive K_{Ca} channels (McManus *et al.*, 1993).

During labour, the forceful contractions of the uterus can occlude its blood supply, which may lead to hypoxia. Hypoxia can reduce and even abolish uterine force in both isolated rat and human uterus (Heaton *et al.*, 1993). Thus, hypoxia may contribute to uterine dystocia (inadequate uterine contraction during labour), the cause of which remains largely unknown and which often results in emergency Caesarean delivery. In a model of hypoxia involving cyanide administration to inhibit uterine force, glibenclamide reduces the K^{+} efflux produced by cyanide (Heaton *et al.*, 1993), thus implicating K_{ATP} channels in this response.

More work is needed to increase our understanding of the K^{+} channels in the myometrium; how they may differ from those in other tissue types and how they may be involved in different pathologic states.

In the treatment of impotence, vasodilators, injected locally, are commonly used. The established vascular smooth muscle relaxant properties of K_{ATP} COs (see Section 4.1.2) suggest that these drugs may offer an alternative to current treatment. Cromakalim and pinacidil increased $^{86}Rb^{+}$ efflux and inhibited spontaneous contractile activity, electrically and noradrenaline-induced contractions in rabbit isolated cavernosal tissue (Holmquist *et al.*, 1990a). Similar results were obtained for pinacidil in human isolated cavernosum (Holmquist *et al.*, 1990b). In addition, pinacidil increased whole cell K^{+} current in human cultured corporal (corpus cavernosum) SMC (Christ *et al.*, 1993). Recently, cromakalim was reported to increase intracavernous pressure in a Simian monkey model, resulting in an erectile response of the penis (Tripp-Rocha *et al.*, 1994). Minoxidil was more effective in facilitating erection and produced fewer side effects than nitroglycerin, when used to treat organic impotence in men (Cavallini, 1991).

4.4. Urinary Bladder

Bladder hyperactivity, secondary to bladder hypertrophy or partial outflow obstruction resulting in urinary incontinence, is common, and the existing therapeutic regimens are often ineffective or poorly tolerated (Wein, 1991). Cromakalim and pinacidil relax urinary bladder smooth muscle, indicating potential in the treatment of urinary incontinence (Andersson *et al.*, 1988; Malmgren *et al.*, 1989). In isolated detrusor muscle preparations from human unstable bladder (due to urinary outflow obstruction), cromakalim inhibited elevated basal tone and spontaneous contractile activity (Foster *et al.*, 1989). Similar effects were also observed with pinacidil (Fivaev *et al.*, 1989). K_{ATP} COs demonstrated inhibitory effects, in both human and animal tissues, on myogenic activ-

ity and contractions to a variety of spasmogens (Longman and Hamilton, 1992). The ability of K_{ATP} COs to inhibit electrically induced contractions of urinary bladder tissues have been variable and may be related to the degree of depolarization produced by neuronal stimulation in different models.

In bladder tissues from a variety of species, K_{ATP} COs (e.g., BRL 38227, pinacidil and RP 49356) increase $^{42}K^{+}$ efflux. However, as found in other smooth muscle systems, the concentrations of drug required are higher than those inhibiting myogenic activity (Longman and Hamilton, 1992). The relaxant activity and enhancement of $^{42}K^{+}$ efflux due to K_{ATP} COs are sensitive to glibenclamide.

Evaluation of K_{ATP} COs on *in vivo* urinary bladder models have been limited due to the vascular effects of the drugs. A comparison of the effects of BRL 38227, pinacidil, Ro 31-6930, RP-49356, and S 0121 did not reveal selectivity for the rat detrusor muscle over portal vein (Edwards *et al.*, 1991). A series of novel K_{ATP} COs (for example, compound ZD6169) recently have been proposed that act selectively on the urinary bladder smooth muscle, without producing significant cardiovascular effects following oral administration to rats (Grant *et al.*, 1994).

The observations in animal models have yet to be supported by clinical trials, as results of initial studies in humans were disappointing. Data from a study with pinacidil in patients with bladder hyperactivity and bladder outflow obstruction (secondary to prostatic hyperplasia) failed to demonstrate any clear improvement in bladder function (Hedlund *et al.*, 1991). Levrocromakalim (BRL 38227) increased the duration of bladder contraction, but was without effect on other urodynamic parameters in patients with high spinal cord lesions (Komersova *et al.*, 1995). Hypotensive responses during this study led the authors to suggest that higher doses of the drug could only be evaluated if administered intravesically.

The relaxant effects of cromakalim and S 0121 (benzopyran K_{ATP} CO) in isolated ureter from rabbit and humans (Klaus *et al.*, 1990) suggest a benefit for K_{ATP} COs in the treatment of kidney stones by aiding their passage along the ureter (Englert *et al.*, 1988). In studies on the guinea-pig ureter, CGRP (see Section 2.3) appears to be an endogenous KCO (Santicioli and Maggi, 1994). The role of CGRP and the involvement of K^{+} channels on human ureter still requires investigation.

4.5. Gastrointestinal Tract

Spontaneous slow-wave contractile activity and/or contractile responses to spasmogens in a variety of gastrointestinal tissues (e.g., taenia caeci, ileum, colon, muscle myenteric plexus, oesophageal, stomach) have been shown to be inhibited by K_{ATP} COs (Longman and Hamilton, 1992). These effects were associated with K^{+} efflux and hyperpolarization that involved a glibenclamide-sensitive mechanism. The animal data suggest that K_{ATP} COs may have utility in conditions associated with disturbances in gastrointestinal motil-

ity, such as irritable bowel syndrome, especially because kinetically slow K^+ channels, carrying outward current, may be responsible for gastrointestinal slow-wave activity (Benham and Bolton, 1983). Interestingly, the evaluation of such drugs in other clinical conditions has not revealed an incidence of adverse side effects on the gastrointestinal tract such as constipation; however, this would be influenced by the site of adsorption of these agents. To gain a full appreciation of potential therapeutic benefits, $KCOs$ are required that are not removed from the gastrointestinal tract.

Diazoxide, like morphine, has a protective effect on ethanol-induced gastric lesions; an effect proposed by Bhounsule *et al.* (1992) to involve $KATP$ channels. As described in Section 4.6.2, opioid receptors are associated with K^+ channels, and glibenclamide is reported to antagonise morphine analgesia (Orano *et al.*, 1990). In the model of gastric lesions, results obtained with glibenclamide were complicated by its action on prostaglandin production in this tissue (Bhounsule *et al.*, 1992).

Neurotensin inhibits contractions in rat and canine ileal smooth muscle by opening opamin-sensitive K_C channels (Alliercher *et al.*, 1992; Christinck *et al.*, 1992). Therefore, SK_CCOs , like $KATPCOs$, may have utility in conditions associated with disturbances in gastrointestinal motility.

4.6. Nervous System

4.6.1. Peripheral. The $KATPCOs$ interfere with neurotransmission in peripheral parasympathetic neurones in the airways and the gastrointestinal tract (Longman and Hemilton, 1992; Section 4.2). This has led to suggestions of a presynaptic site of action, whereby the $KATPCOs$ control the release of neurotransmitter. In contrast, cromakalim, nicotinic, and pinacidil failed to exert an inhibitory effect on noradrenaline release in rat isolated mesenteric artery (Fabiani and Story, 1994). The presynaptic inhibitory role of $KATPCOs$, therefore, appears to be selective for parasympathetic (cholinergic) innervation.

Interestingly, cromakalim and pinacidil inhibited nicotinic acetylcholine receptor mediated and voltage-dependent catecholamine secretion from bovine adrenal chromaffin cells (Masada *et al.*, 1994). Thus, $KATPCO$ -sensitive K^+ channels could be involved in regulation of catecholamine secretion mainly indirectly through $VOCs$.

4.6.2. Central. Potassium channels play a pivotal role in the control of neuronal excitability, action potential, and neurotransmitter release within the CNS (Hille, 1984; Cox, 1990). Activation of a variety of receptors [e.g., opioid, 5-HT₁, somatostatin, α -adrenoceptors] by the appropriate neurotransmitter alters the flux of K^+ ions from neurones (North, 1989). Because of this role in normal CNS physiology, derangements in the function of K^+ channels may underlie several CNS diseases. Studies of the distribution of binding sites for the three ligands, [¹²⁵I]iodoglyburide, [¹²⁵I]apamin and [¹²⁵I]charybdoxin, as markers for $KATP$, SK_C , and BK_C (or voltage-gated K^+) channels, respectively, was dramatically different in the rat brain (Giehlert and

Gackheimer, 1993). These data indicate that pharmacological modulation of these K^+ channel subtypes should result in distinctly different effects on brain function. [¹²⁵I]iodoglyburide binding exhibits a very broad distribution in brain, being found in a majority of brain regions. The globus pallidus and the zona reticularis of the substantia nigra (involved in movement coordination) contain the highest density of binding. Openers of $KATP$ channels have been suggested to exert a protective effect on ischemic tissue by reversing the ischemia-induced depolarization (see below). Therefore, $KATPCOs$ may find utility as neuroprotective drugs, and the broad distribution of these channels would indicate that these drugs would have an effect on most neuronal populations in the brain. Although $KATPCO$ ligands, [¹²⁵I]PI075 and [¹²⁵I]cromakalim, are available, a profile of the binding sites distribution within the CNS is not published as yet. [¹²⁵I]Apamin binding suggested that SK_C channels are associated primarily with cell bodies and dendritic spines, rather than nonneuronal elements. Localization in the cerebral cortex and hippocampus would suggest that this channel may have a role in the processing of memory. Interestingly, a loss of [¹²⁵I]apamin binding sites has been reported in the subiculum and CA₁ neurones of hippocampal tissue obtained from postmortem Alzheimer's disease patients (Ikeda *et al.*, 1991). A reasonable hypothesis is that openers of the SK_C channel may confer a neuroprotective effect; however, development of appropriate chemical molecules and pharmacological tools is awaited. The highest levels of [¹²⁵I]charybdoxin binding sites were found in the white matter-containing regions, such as the lateral olfactory tract and fasciculus reticulospinal. This suggests that the charybdoxin-sensitive K^+ channel is present on axons and may modulate nerve conduction in these regions.

Studies *in vitro* and in animal models indicate the potential clinical utility of $KATPCOs$ for diseases of the CNS.

During anoxic conditions, neuronal depolarization is due, at least, to the release of large concentrations of excitatory amino acids, such as glutamate, which may be involved in long-term ischemia-induced damage in the brain (Miller, 1990). *In vitro* experiments, diazoxide and somatostatin were shown to prevent anoxia-induced depolarization of CA3 hippocampal neurones following the opening of K^+ channels (Ben-Ari *et al.*, 1990); these effects were inhibited by pretreatment with glyburide. The authors proposed that $KATPCOs$ may prevent anoxia-induced damage to hippocampal neurones by inhibiting the release of excitatory amino acids. This suggestion is supported by the finding that levcromakalim and RP 52891 blocked ischemia-induced glutamate release in rat hippocampal slices (Zini *et al.*, 1993). In addition, the $KATPCOs$ levcromakalim, nicotinic, and pinacidil blocked ischemia-induced expression of the genes *c-fos* and *c-jun* and of the mRNAs for 70-kDa heat-shock protein and the form of the amyloid β -protein precursor, including the Kurumi-type protease inhibitor domain in rat hippocampus (Heurteaux *et al.*, 1993).

The movement disorders associated with Parkinson's dis-

case are due to a selective loss of dopaminergic neurons in the substantia nigra. The highest density of K_{ATP} s, as judged by autoradiographic studies with [125 I]-iodoglyburide, in the brain is found in the substantia nigra (Gehlert and Gackenheimer, 1993). Sulphonylurea binding studies, however, provide essentially only indirect evidence of K_{ATP} s. Sulphonylureas or increasing extracellular glucose increase the release of [14 C]GABA from the substantia nigra, effects that are inhibited by K_{ATP} COs (Schmid-Antomarchi *et al.*, 1990). GABAergic pathways to other CNS structures (e.g., raphe nuclei) are also modified by K_{ATP} COs. Schmid-Antomarchi *et al.* (1990) noted that the order of potency of the K_{ATP} COs (levromakalim > nicotrandil > cromakalim > diazoxide > pinacidil) was found to be different from that in either the pancreatic β -cell or in smooth muscle, possibly indicating a difference between the target K^+ channel in this brain region and that in other tissues. The K_{ATP} channel in neuronal tissue is not the classical (Type I) channel found in pancreatic β -cells or heart, but a large-conductance nonrectifying version (Type 2; Ashcroft and Ashcroft, 1990).

The genesis and propagation of nonphysiologic electrical impulses are the hallmark of epilepsy. Thus, the hyperpolarization (and restraining) of excitable cells through the opening of K^+ channels could demonstrate therapeutic benefit in this setting. The K_{ATP} COs cromakalim and RP 52091 (isprukalim) reduced seizures in genetically epileptic rats (Gardiolto *et al.*, 1989) and pentylenetetrazole-induced seizures in mice are blocked by intracerebroventricular administration of cromakalim, but not pinacidil (Del Pozo *et al.*, 1993). In a diazepam-induced model of tonic-clonic seizures in the rat, cromakalim completely inhibited both EEG and behavioural seizures, and picrobarbitone only prevented behavioural activity (Popoli *et al.*, 1991). K_{ATP} COs counteract anoxic hyperexcitability, but not 4-aminopyridine-induced epileptiform activity in the rat hippocampal slice (Marta *et al.*, 1994), suggesting such drugs might be useful in the treatment of seizures occurring in the setting of status epilepticus or cerebrovascular disease. High concentrations of cromakalim inhibit electrically induced epileptiform discharges in guinea-pig hippocampus (Alzheimer and von Bruggencate, 1988). Finally, the antiepileptic drug carbamazepine increases potassium currents in rat cortical neurons (Torres *et al.*, 1990). However, whether or not this property is linked to the pharmacological actions of the drug requires further investigation. A similar property (K^+ channel opening), which may be of clinical importance, has been reported to be exhibited by oxcarbazepine (Mullen *et al.*, 1994).

Opioids exert their analgesic effects by binding to opiate receptors, which leads to opening of K^+ channels and neuronal hyperpolarization (North, 1989). Morphine-induced antinociception in mice tail-flick tests is mediated by the opening of K_{ATP} channels (Ocana *et al.*, 1993). These observations would suggest a potential role for K_{ATP} COs as analgesics. Intrathecal administration of the K_{ATP} COs BRL-38227, minoxidil, and diazoxide produced antinociception in the tail-flick test in mice (Welch and Dunlow, 1993). The

K_{ATP} COs were not cross-tolerant to the effects of morphine in this model. This led to the suggestion that the K_{ATP} COs and opioids probably do not act on a common site, but could have a common second messenger (Welch and Dunlow, 1993). A dose-dependent increase in the effects of morphine on the hot-plate and tail-flick tests were obtained following i.v. administration of pinacidil to rats (Vergoni *et al.*, 1992).

Benzo[pyran] derivatives of cromakalim (e.g., SR 46142A) have been claimed to exert antidepressant activity in animal tests (improve swimming performance in mice) in the absence of a cardiovascular effect (Garcia *et al.*, 1990). Similar results were observed with a series of arandochromans (Poucellet, 1990). This would suggest that modifications of the benzo[pyran] nucleus of cromakalim allows tissue selectivity (CNS over vascular smooth muscle) to be achieved. Whether or not the antidepressant effects of SR 46142A (and related compounds) are mediated by modulation of K^+ channels remains to be confirmed.

These studies are encouraging, but neuronal tissue-selective agents that cross the blood-brain barrier are needed to realize the potential clinical applications of K_{ATP} COs in CNS disorders. In addition, the promiscuity of K_{ATP} COs invites adverse CNS effects; for example, cromakalim, like d-amphetamine, enhances spontaneous locomotor activity in the rat by a glubenzamide-sensitive mechanism (Analfic *et al.*, 1992).

BK_{Ca} channels also play a role in the function of neuronal cells (Hille, 1984; Cook, 1990). The recent identification of compounds (e.g., NS 1619, see Section 2.2) that activate BK_{Ca} channels will allow determination of their utility as potential therapeutic agents. In rat cerebellar granule cells, NS-004 activates a 187 pS Ca -dependent K^+ channel that is blocked by charybotoxin (Olesen *et al.*, 1994a). NS 1619 activates the BK_{Ca} , but not K_{ATP} channels, in membrane patches isolated from rat ventromedial hypothalamic neurones (Sellers and Ashford, 1994). Although K_{ATP} channels are present in rat ventromedial hypothalamic neurones, BRL 38227, cromakalim, and pinacidil all failed to evoke an effect on these channels in excised membrane patches (Sellers *et al.*, 1992). The glucose sensitive cells in the ventromedial hypothalamus are involved in the control of appetite and are often regarded as the satiety center (Morley, 1980; Blundell, 1992). Thus, activation of BK_{Ca} channels would decrease hypothalamic firing and reduce the sensation of satiety.

4.7. Skeletal Muscle

In skeletal muscle, K_{ATP} channel activity has been shown to increase upon intracellular acidification (Davies, 1990). Falls in intracellular pH reduce the inhibitory effect of ATP on K^+ channels in frog skeletal muscle. This could mean that during increased muscle exercise and consequent lowering of pH, K_{ATP} channel-induced hyperpolarization could compensate for a decrease in electrical excitability and prevent spontaneous contractions from occurring.

The K_{ATP} COs, cromakalim, pinacidil, and RP-49356,

increased opening time of a glibenclamide-sensitive K^+ channel in mouse skeletal muscle (Weik and Neumcke, 1990). Dioxazole activated K_{ATP} channels in smooth muscle and pancreatic cells, but had very little effect in skeletal muscle, even at very high concentrations (Weik and Neumcke, 1990). Interestingly, this sulphonamide (thus, structurally related to the sulphothiazes) inhibited ATP-sensitive channels in ventricular muscle cells (Faivre and Findlay, 1989). Thus, the different effects of dioxazole suggest that the K^+ channels in mouse skeletal muscle resemble those in cardiac cells more than those in smooth muscle and pancreatic β -cells (Lawson and Hicks, 1993). Due to diverse effects of pinacidil in mouse skeletal muscle, a model of the K_{ATP} channel was proposed with a binding site for the $K_{ATP}CO$ and two sites for nucleotides, one activating and one inhibitory (ATP or ADP can occupy either site; Fiehl and Neumcke, 1994). Pinacidil activates the channel and displaces the blocker from the inhibitory site, only if the activating site is occupied.

Cromakalim enhanced K^+ efflux in human skeletal fibres, an effect that was blocked by tolbutamide (Spuler *et al.*, 1989), suggesting $K_{ATP}COs$ could have some role in the treatment of pathological muscle fatigue or paralysis resulting from excessive membrane depolarization. Interestingly, cromakalim, pinacidil, and RP-49356 evoked larger hyperpolarizations in skeletal muscle fibres from patients with myotonic dystrophy or hypokalaemic periodic paralysis than in those from normal volunteers (Spuler *et al.*, 1989; Quasthoff *et al.*, 1989; Giale *et al.*, 1990). $K_{ATP}COs$, however, increased open probability of an ATP-sensitive and an ATP-insensitive K^+ channel in human skeletal muscle (Quasthoff *et al.*, 1990); the effect of the $K_{ATP}CO$ on both channels was blocked by glibenclamide.

Ischaemia-induced damage in a rat skeletal muscle, like cardiac (Section 4.1.1) and neuronal (Section 4.6.2) ischaemia, has been shown to be prevented by cromakalim (Haiton *et al.*, 1991). This result, and the observation that cromakalim restores the membrane potential of depolarized human skeletal muscle fibres (Spuler *et al.*, 1989), indicate that $K_{ATP}COs$ may be useful for the treatment of peripheral vascular disease (see Section 4.1.1). In rat skeletal muscle, Angersbach and Nicholson (1988) demonstrated that $K_{ATP}COs$, but not Ca^{2+} antagonists or hydralazine, selectively increase blood flow to collateral vessels in a previously ischaemic limb. Weikewich *et al.* (1993), however, suggested that $K_{ATP}COs$ would not be beneficial in treatment of skeletal muscle ischaemia *in vivo*, but may be useful in preserving skeletal muscle function in cases of ischaemia followed by reperfusion.

Studies in human muscle have implicated SK_{Ca} (tapamin-sensitive) channels in the condition myotonic muscular dystrophy, characterized by muscle stiffness (Renaud *et al.*, 1986). Although K^+ channel subtypes other than K_{ATP} channels exist in skeletal muscle (SK_{Ca} , BK_{Ca} , delayed rectifier K^+ channel; Wareham, 1992), however, selective openers are awaited to determine their role and the therapeutic potential of activation.

4.8. Hair Growth

The occurrence of hypertrichosis during antihypertensive treatment with minoxidil (Campese, 1981) led to the subsequent evaluation of the drug (applied topically) to enhance hair regrowth in areas of baldness (Chisold and Heel, 1987). Topically administered minoxidil enhances hair growth in certain forms of male pattern baldness. Although this effect has been suggested to involve K^+ channels, hypertrichosis occurs in 80–100% of minoxidil-treated patients, but only 2–13% of pinacidil-treated patients. There is no evidence that cromakalim, nicorandil, or RP-49356 stimulate hair growth. In SMC, minoxidil (sulphate) has been suggested to open a K^+ channel that is not recognized by other $K_{ATP}COs$ (see Section 2.1). $K_{ATP}COs$ stimulate DNA synthesis in mouse epidermal keratinocyte and whole hair follicle cultures (Herman *et al.*, 1993). In cultured whisker follicles, minoxidil, but not P1075 (pinacidil analog), preserved the root sheath; however, both drugs stimulated cysteine incorporation in follicles (Walden *et al.*, 1993). The root sheath may be the target for minoxidil; thus, stimulating hair growth through a direct effect on the hair follicle.

4.9. Intraocular Pressure

Repeated topical application of pinacidil, cromakalim, or nicorandil lowers intraocular pressure (IOP) of rabbits, suggesting a potential benefit of $K_{ATP}COs$ in eye disorders such as glaucoma (Godtfredsen, 1989). In an isolated arterially perfused bovine eye preparation, pinacidil caused a sustained decrease in IOP, with no effect on arterial perfusion pressure (Miller and Wilson, 1991); thus, suggesting that relaxation of resistance vessels is not involved in the fall in IOP due to the $K_{ATP}CO$. Whether these effects in the eye may be attributed to enhanced K^+ ion movement and consequent relaxation of smooth muscle or are the result of K^+ channel modulation in epithelial cells requires further investigation.

5. CONCLUSIONS

Synthetic molecules that 'directly' activate K^+ channels have led to a new direction in the pharmacology of ion channels. The identification of the K^+ channel opening property of cromakalim initiated major research efforts in the search for other such agents and in the determination of the specific channel(s) involved. The existence of so many different subtypes of K^+ channels has been an impetus in the search for new molecules that would have different profiles and channel selectivities (e.g., K_{ATP} , BK_{Ca}). The availability of an increasing number of $KCOs$, exogenous and endogenous, should facilitate more detailed study of these channels under both normal and certain pathophysiological conditions. The decrease in the excitability of cells that follows K^+ channel opening offers a broad clinical potential for drugs with this property in a number of pathological conditions. Consequently, therapeutic roles of $KCOs$ can be envisaged in disorders of a wide range of cells; for example, vascular and nonvascular smooth muscle,

cardiac, neuronal, and skeletal cells. Although lack of selectivity of current compounds remains one of the major hurdles in this area, advances in prototype KCOs and our knowledge of K⁺ channel pharmacology is encouraging. Thus, the development of KCOs that will provide positive results in extensive clinical trials to give an appreciation of the full therapeutic potential are eagerly awaited.

Acknowledgments: Dr. E.E. Hicks (Quelined) is thanked for his constructive comments. The authors also gratefully thank Anne Lawson for assistance in the preparation of this article.

References

- Adeagbo, A. S. O. and Triggle, C. R. (1993) Varying extracellular [K⁺]: a functional approach to separating EDHF-related and EDHF-related mechanisms in perfused rat mesenteric arterial bed. *J. Cardiovasc. Pharmacol.* 21: 423-429.
- Adeagbo, A. S. O., Triggle, C. R., and Blegen, D. (1993) Desopthalgic smooth muscle relaxation by dimethylphenylpiperazinium (DMPP) relationship to muscarinic cholinergic receptor antagonism and K⁺ channel activation. *Pharmacol. Commun.* 2: 285-294.
- Amleth-Renore, L. (1988) Piroxicam: history, basic pharmacology and therapeutic implications. *J. Cardiovasc. Pharmacol.* 12: S1-S4.
- Allen, S. L., Boyle, J. P., Corliss, J., Foster, R. W., Morgan, G. P., and Small, R. C. (1986) Electrical and mechanical effects of BRL 14915 in guinea-pig isolated trachea. *Br. J. Pharmacol.* 89: 395-405.
- Altschuld, H. D., Fick, H., Schudtziarra, V., and Classen, M. (1992) Mechanisms of neurotensin-induced inhibition in rat skeletal muscle. *Am. J. Physiol.* 263: C767-C774.
- Altschuld, C. and Ten Bruggencate, G. (1985) Actions of BRL 14915 (fen-nakalim) upon convulsive discharges in guinea-pig hippocampal slices. *Naunyn-Schmiedeberg Arch. Pharmacol.* 337: 429-434.
- Amis, M., Heurteaux, C., Nicod, P., and Lanthier, M. (1992) Behavioural effects of modulators of ATP-sensitive K⁺ channels in the rat dorsal pallidum. *Eur. J. Pharmacol.* 217: 71-77.
- Arpaci, O. (1977) Flatus that kill. *World Health* 26: 28-33.
- Andersson, K.E., Andersson, P. G., Fossius, M., Hedlund, H., Malmgren, A., and Sjogren, C. (1988) Effects of piroxicam on bladder muscle. *Drugs* 36: 41-49.
- Argersinger, D. and Nicholson, E. (1988) Enhancement of muscle blood cell flux and pO₂ by cromakalim (BRL 14915) and other compounds enhancing membrane K⁺ conductance, but not by Ca²⁺ antagonists or hyaluronate, in an animal model of occlusive arterial disease. *Naunyn-Schmiedeberg Arch. Pharmacol.* 337: 341-346.
- Arnsperger, C. and Di Diego, J. M. (1992) Role of K⁺ channel activation in cardiac electrophysiology and arrhythmias. *Circulation* 85: 1627-1629.
- Atwell, K., Chen, C., Perez, G. J., Pateroyan, N., McDougall, J. K., Munga, M., Sanborn, B. H., Sreima, E., and Tori, L. (1993) Calcium activated K⁺ channels as modulators of human myocardial contractile activity. *Am. J. Physiol.* 265: C976-C985.
- Arch, J. R. S., Taylor, S. G., and Buckle, D. R. (1992) Potassium channel activators and the treatment of asthma. *Current Drugs Potassium Channel Modulators* K.C. M. B139-KC. M. B150.
- Arch, J. R. S., Bowring, J. L., and Buckle, D. R. (1994) Evaluation of the novel potassium channel activator BRL 15534 as an inhaled bronchodilator in guinea-pig and rats: comparison with fenoterol and salbutamol. *Pulm. Pharmacol.* 7: 121-128.
- Aronson, J. K. (1992) Potassium channels in nervous tissue. *Biochem. Pharmacol.* 43: 11-14.
- Arundel, H. and Schild, H. O. (1959) Some quantitative uses of drug antagonists. *Br. J. Pharmacol. Chemother.* 16: 48-58.
- Ascher, S. H. and Ascher, F. M. (1990) Properties and functions of ATP-sensitive K⁺ channels. *Crit. Rev. Toxicol.* 2: 197-219.
- Awai, K. S. (1992) Modulation of potassium channels by organic molecules. *Med. Res. Rev.* 12: 569-591.
- Awai, K. S. (1994) Pharmacology and structure activity relationships for K_{ATP} modulators—use selective K_{ATP} openers. *J. Cardiovasc. Pharmacol.* 24: S12-S17.
- Audampach, J. A., Maruyama, M., Caverio, L., and Gross, G. J. (1992) Pharmacological evidence for a role of ATP-dependent potassium channels in myocardial stunning. *Circulation* 86: 311-319.
- Barnes, P. J., Beranek, J. N., and Belvisi, M. G. (1991) Neuropeptides in the respiratory tract. *Am. Rev. Respir. Dis.* 144: 1187-1198.
- Beech, D. J., Zhang, H., Nakao, K., and Bolton, T. B. (1993a) K⁺ channel activation by nucleotide diphosphates and its inhibition by glibenclamide in vascular smooth muscle. *Br. J. Pharmacol.* 110: 573-582.
- Beech, D. J., Zhang, H., Nakao, K., and Bolton, T. B. (1993b) Single channel and whole cell K⁺ currents evoked by 10-nM glibenclamide in smooth muscle cells from the rabbit portal vein. *Br. J. Pharmacol.* 110: 583-590.
- Ben-Ari, Y. (1990) GABA and glibenclamide modulate the ionic release of glutamate in rat CA1 hippocampal neurons. *Eur. J. Neurosci.* 2: 62-68.
- Ben-Ari, Y., Kresche, K., and Orde, P. (1990) Activators of ATP-sensitive K⁺ channels reduce anoxic depolarisation in CA1 hippocampal neurons. *Neuroscience* 37: 55-61.
- Benham, C. S. and Bolton, T. B. (1983) Patch-clamp studies of slow potential sensitive potassium channels in longitudinal smooth muscle cells of rabbit jejunum. *J. Physiol. (Lond.)* 345: 469-486.
- Berry, J. L., Elliot, K. R. F., Foster, R. W., Green, K. A., Murray, M. A., and Small, R. C. (1991) Mechanisms, biochemical and electrophysiological studies of BRL 14915 and cromakalim in guinea-pig trachealis muscle. *Pulm. Pharmacol.* 4: 91-98.
- Brown, J. A. and Bayliss, J. E. (1986) Noradrenergic neural vasodilator mechanisms. *Circ. Res.* 60: 309-326.
- Brown, S. A., Dinitz, D'Souza, R. S., and Sharma, V. G. (1992) Protective effect of morphine on ethanol-induced gastric lesions in rats: are ATP-dependent potassium channels involved? *Arch. Int. Pharmacodyn.* 318: 126-133.
- Burn, K. and Hermsmeider, K. (1994) Glyburide actions on the dihydropyridine-sensitive Ca²⁺ channel in rat vascular muscle. *J. Vasc. Res.* 31: 256-264.
- Black, J. L., Armour, C. L., Johnson, P. R. A., Alouani, L. A., and Barnes, P. J. (1990) The action of a potassium channel activator, BRL 16227 (lenakalim), on human airway smooth muscle. *Am. Rev. Respir. Dis.* 142: 1384-1389.
- Black, S. C. and Luchini, B. R. (1994) Potassium channel openers are likely to be proarrhythmic in the diseased human heart. *Cardiovasc. Res.* 28: 923-924.
- Blatt, A. L. and Magley, K. L. (1987) Calcium-activated potassium channels. *Trends Neurosci.* 10: 463-467.
- Blundell, J. (1992) Pharmacological approaches to appetite suppression. *Trends Pharmacol. Sci.* 12: 147-152.
- Boeyens, J. M. and Rambert, J. (1990) Potassium channels and endothelial prostacyclin production: in importance of Prostacyclin and Potassium Channels for the Regulation of Vascular Tone, pp. 31-36, Frolich, J. C., and Fostermann, K. (eds.) Zuckerswerdt, Munich.

- Bolton, V. M., Naphis, S., Palacina, J. J., Pagaris, P. J. and Cohen, R. A. (1994) Nitric oxide directly activates calcium-dependent potassium channels in vascular smooth muscle. *Nature* 368: 850-853.
- Bowring, N. E., Buckle, D. R., Clarke, G. D., Taylor, J. F. and Arch, J. R. S. (1991) Evaluation of the potassium channel activator BRL 38227 as an inhaled bronchodilator in the guinea-pig trachea with rufledipine and salbutamol. *Pulm. Pharmacol.* 4: 99-105.
- Bowring, N. E., Arch, J. R. S., Buckle, D. R., Clarke, G. D. and Taylor, J. F. (1993) Comparison of the smooth relaxant and hypotensive potencies of the potassium channel activators BRL 38227 and levcromakalim (BRL 38227) in vivo in guinea-pig and rats. *Br. J. Pharmacol.* 109: 1133-1139.
- Beay, K. M., Newgreen, D. T., Small, R. C., Southerton, J. S., Taylor, N. G., Wen, S. W. and Weston, A. H. (1987) Evidence that the mechanism of the inhibitory action of pinacidil in rat and guinea-pig smooth muscle differs from that of glyceryl trinitrate. *Br. J. Pharmacol.* 91: 421-429.
- Bray, V. M., Weston, A. H., Duty, S., Newgreen, D. T., Longmore, J., Edwards, G. and Brown, T. J. (1991) Differences between the effects of cromakalim and rufledipine on agonist-induced responses in rabbit aorta. *Br. J. Pharmacol.* 102: 337-344.
- Bulbring, E. and Yama, T. (1987) Catecholamine action on smooth muscle. *Pharmacol. Rev.* 39: 49-96.
- Burka, J. F., Berry, J. L., Foster, R. W., Small, R. C. and West, A. J. (1991) Effects of cromakalim on neurally-mediated responses of guinea-pig tracheal smooth muscle. *Br. J. Pharmacol.* 104: 263-269.
- Calder, J. A., Schachter, M. and Sever, P. S. (1994) Potassium channel opening properties of thiazide diuretics in isolated guinea-pig resistance arteries. *J. Cardiovasc. Pharmacol.* 24: 158-164.
- Campese, V. M. (1981) Minoxidil—a review of its pharmacological properties and therapeutic uses. *Drugs* 22: 257-278.
- Carlson, J. E., Kandel, T., Jensen, H. E., Tang, M. and Trap-Jensen, J. (1983) Pinacidil: a new vasodilator: pharmacokinetics of a new retarded release tablet in essential hypertension. *Eur. J. Pharmacol.* 75: 557-561.
- Carlson, J. E., Kandel, T., Lund, J. O., Mohab, A. and Trap-Jensen, J. (1985) Acute haemodynamic effects of pinacidil and hydralazine in essential hypertension. *Clin. Pharmacol. Ther.* 37: 251-259.
- Carlier, L., Abramson, C., Drews, L. and Duker, G. (1991) Arrhythmogenic effects of potassium channel openers in rhythmic abnormalities related to delayed repolarization. *Circulation* 85: 1491-1506.
- Casade, N. A., Haylett, D. G. and Jenkinson, D. H. (1989) Toxins in the characterization of potassium channels. *Trends Neurosci.* 12: 39-45.
- Cavallini, G. (1991) Minoxidil versus nitroglycerine: a prospective double-blind controlled trial in transcutaneous oxygen saturation for organic impotence. *J. Urol.* 146: 50-53.
- Caveys, I. and Permeteux, J. (1994) ATP potassium channel openers are of potential benefit in ischemic heart disease. *Cardiovasc. Res.* 28: 12-33.
- Carver, L., Lawson, K. and Pratt, J. (1982) An investigation of K⁺-induced relaxation of the rat aorta suspended in a K⁺ deficient solution. *Br. J. Pharmacol. Proc. Suppl.* 91: 455P.
- Chapman, I. D., Manzoni, L. and Morley, J. (1991) SLZ PCO 400 and cromakalim suppressed development of airway hyperactivity at concentrations which do not induce overt smooth muscle relaxation. *Agents Actions* 34 (Suppl.): 53-62.
- Chen, G. and Cheung, D. W. (1992) Characterization of acetylcholine-induced membrane hyperpolarization in endothelial cells. *Circ. Res.* 70: 257-263.
- Chen, G. and Suzuki, H. (1989) Some electrical properties of the endothelium-dependent hyperpolarization recorded from rat arterial smooth muscle cells. *J. Physiol. (Lond)* 410: 91-106.
- Chen, J. M. S., Hollingsworth, M., Hughes, S. J., Piper, I. T. and Marelli, M. J. A. (1993) Inhibition of constriction of the isolated human myocardium by potassium channel openers. *Am. J. Obstet. Gynecol.* 168: 953-960.
- Chi, L., U'Prichard, A. C. D. and Lucchesia, B. R. (1990) Prothrombotic actions of pinacidil in a conscious canine model of sudden coronary death. *J. Cardiovasc. Pharmacol.* 15: 452-464.
- Chioara, L. C., Twort, C. H. C. and Waid, J. P. T. (1992) Direct action of BRL 38227 and glibenclamide on intracellular calcium stores in cultured airway smooth muscle of rabbit. *Br. J. Pharmacol.* 105: 259-260.
- Christ, G. J., Spray, D. C. and Brink, P. R. (1993) Characterization of K⁺ currents in cultured human corneal smooth muscle cells. *J. Neurosci.* 13: 319-328.
- Christinck, F., Daniel, E. and Knochel, J. E. T. (1992) Inhibitory and excitatory mechanisms of neurotensin action in canine intestinal vascular muscle in vitro. *Can. J. Physiol. Pharmacol.* 70: 1423-1431.
- Clossold, S. F. and Heel, R. C. (1987) Topical minoxidil: A preliminary review of its pharmacodynamic properties and therapeutic efficacy in alopecia areata and alopecia androgenetica. *Drugs* 33: 107-122.
- Coetzee, W. A. (1992) ATP-sensitive potassium channels and myocardial ischemia: why do they open? *Cardiovasc. Drugs Ther.* 6: 201-208.
- Coldwell, M. C. and Howlett, D. R. (1988) Potassium efflux enhancement by cromakalim (BRL 34915) in rabbit mesenteric artery: an indirect effect independent of calcium. *Biochem. Pharmacol.* 37: 4105-4110.
- Cook, D. L. and Hales, C. N. (1984) Intracellular ATP directly blocks K⁺ channels in pancreatic β -cells. *Nature* 311: 271-273.
- Cook, N. S. (1990) Potassium Channels: Structure, Classification, Function and Therapeutic Potential. Ellis Horwood Ltd, Chichester.
- Cook, N. S. and Chapman, I. D. (1993) Therapeutic potential of potassium channel openers in peripheral vascular disease and asthma. *Cardiovasc. Drugs Ther.* 7: 355-363.
- Cook, S. J., Arcieri, K., Martin, A., Bucher, K. H., Forder, J. R., Moller, T., Miller, A. J., Elliott, K. B. F., Foster, R. W. and Small, R. C. (1995) Further analysis of the mechanisms underlying the tracheal relaxant action of SCA 40. *Br. J. Pharmacol.* 114: 143-151.
- Cooke, J. F., Rasmach, E., Andon, N. A., Loscalzo, J. and Dzau, V. J. (1991) Flow activates an endothelial potassium channel to release an endogenous nitric oxide. *J. Clin. Invest.* 88: 1663-1671.
- Crepel, V., Krnjevic, K. and Ben-Ari, Y. (1993) Sulphonylureas reduce the slowly inactivating D-type outward current in rat hippocampal neurons. *J. Physiol. (Lond)* 466: 59-54.
- Cridde, D. N. and Soares de Moura, B. (1995) Inhibitory effects of Rb⁺ on the responses to levcromakalim and P1600 in the isolated human myocardium. *Eur. J. Pharmacol.* 272: 293-296.
- D'Alonzo, A. J. and Grover, G. J. (1994) Potassium channel openers are unlikely to be proarrhythmic in the diseased human heart. *Cardiovasc. Res.* 28: 924-925.
- Davies, N. W. (1990) Modulation of ATP-sensitive K⁺ channels in skeletal muscle by intracellular protons. *Nature* 343: 375-377.

- De La Couraie, J. E., Eledjorn, J. J., Bruelle, P., Peray, P. A., Bassoul, B. P., Gagnon, J. P. and Sassiou, A. (1993) Electrophysiological and arrhythmogenic effects of the potassium channel agonist BRL-35227 in anesthetized dogs. *J. Cardiovasc. Pharmacol.* 22: 777-782.
- Del Pozo, E., Barrios, M. and Banyasz, J. M. (1990) Effects of potassium channel openers on pentyltetracetic acid-induced seizures in mice. *Pharmacol. Toxicol.* 67: 182-184.
- De Weille, J., Schmid-Armstrong, H., Fosset, M. and Lazdunski, M. (1988) ATP-sensitive K^+ channels that are blocked by hypoglycemia-inducing sulfonylureas in insulin-secreting cells are activated by galin, a hyperglycemia-inducing hormone. *Proc. Natl. Acad. Sci. USA* 85: 1312-1316.
- De Weille, J., Schmid-Armstrong, H., Fosset, M. and Lazdunski, M. (1989) Regulation of ATP-sensitive K^+ channels in insulinoma cells: activation by somatostatin and protein kinase C and the role of cAMP. *Proc. Natl. Acad. Sci. USA* 86: 2971-2975.
- Dreyer, E. (1990) Peptide toxins and potassium channels. *Rev. Physiol. Biochem. Pharmacol.* 115: 93-136.
- Drieu, L., Rochette, C., Richard, V., Dubois-Randé, J. L., Roupier, E., Giudicelli, J. F., Hirtling, L. and Bendaux, A. (1992) Potassium channel openers dilate large epicardial coronary arteries in conscious dogs by an indirect, endothelium-dependent mechanism. *J. Pharmacol. Exp. Ther.* 263: 1091-1099.
- Duty, S. and Weston, A. H. (1992) Characteristics of the contractile response of rabbit aorta produced by cromakalim in calcium-free solution. *Br. J. Pharmacol.* 107: 1198-1204.
- Eckl, K. M. and Greb, W. H. (1987) Potassium channel activation in the vascular smooth muscle cell membrane - a new mechanism of action in antihypertensive treatment - early clinical results with BRL 34915. *Clin. Exp. Hypertens.* 49: 160.
- Eckman, D. M., Frankovich, J. D. and Keef, K. D. (1992) Comparison of the effects of acetylcholine and BRL-35227 in the guinea-pig coronary artery. *Br. J. Pharmacol.* 106: 9-16.
- Edwards, G. and Weston, A. H. (1990) Structure-activity relationship of K channel openers. *Trends Pharmacol. Sci.* 11: 417-422.
- Edwards, G. and Weston, A. H. (1993) The pharmacology of ATP-sensitive potassium channels. *Annu. Rev. Pharmacol. Toxicol.* 33: 597-637.
- Edwards, G. and Weston, A. H. (1994) K_{ATP} - fact or artefact? New thoughts on the mode of action of the potassium channel openers. *Cardiovasc. Res.* 28: 735-737.
- Edwards, G. and Weston, A. H. (1995) Pharmacology of the potassium channel openers. *Cardiovasc. Drugs Ther.* 9: 185-193.
- Edwards, G., Henshaw, M., Miller, M. and Weston, A. H. (1993) Comparison of the effects of several potassium channel openers on rat bladder and rat portal vein *in vitro*. *Br. J. Pharmacol.* 102: 679-686.
- Edwards, G., Niederste-Hollenberg, A., Schneider, J., Noack, Th. and Weston, A. H. (1994) Ion channel modulation by NS-1619, the putative IK_{ATP} channel opener in vascular smooth muscle. *Br. J. Pharmacol.* 115: 1538-1547.
- Edwards, G., Schneider, J., Niederste-Hollenberg, A., Noack, Th. and Weston, A. H. (1995) Effects of BRL 55834 in rat portal vein and bovine trachea evidence for the induction of a glibenclamide-resistant, ATP-sensitive potassium channel. *Br. J. Pharmacol.* 115: 1027-1037.
- Eliaz, M. (1989a) Glibenclamide is a competitive antagonist of cromakalim, pinacidil and RP 49356 in guinea-pig pulmonary artery. *Eur. J. Pharmacol.* 165: 231-239.
- Eliaz, M. (1999b) Competitive antagonism by glibenclamide of cromakalim inhibition of twitch contractions in rabbit vas deferens. *Eur. J. Pharmacol.* 161: 103-106.
- Englert, C. H., Klaus, E., Lang, H. J., Mann, D. and Scholten, B. (1988) European Patent Application. E. 0227 631.
- Escande, D. and Claverie, L. (1992) K^+ channel openers and 'natural' cardioprotection. *Trends Pharmacol. Sci.* 13: 269-271.
- Fabiani, M. E. and Story, D. F. (1994) Functional effects of cromakalim, nicorandil and pinacidil on noradrenergic transmission in rat isolated mesenteric artery. *J. Auton. Pharmacol.* 14: 87-93.
- Faivre, J. F. and Fridlay, L. (1989) Effects of rubatamide, glibenclamide and diazoxide upon action potentials recorded from rat ventricular muscle. *Biochem. Biophys. Acta* 994: 1-5.
- Fauschou, P., Mikkelsen, K. L., Steffensen, L. and Fratrik, B. (1994) The lack of bronchodilator effect and the short-term safety of cumulative single doses of an inhaled potassium channel opener (bimakalim) in adult patients with mild to moderate bronchial asthma. *Pulm. Pharmacol.* 7: 291-297.
- Foster, C. D., Spakman, M. J., Fujii, K. and Brading, A. F. (1989) The effects of cromakalim in the detrusor muscle of human and pig urinary bladder. *Br. J. Urol.* 63: 284-294.
- Fowles, M., Andersson, K. E. and Hedlund, H. (1989) The action of pinacidil in the isolated human bladder. *J. Urol.* 141: 637-640.
- Fulginiti, J., Cohen, M. M. and Morand, R. S. (1993) Endothelin differentially affects rat gastric longitudinal and circular smooth muscle. *J. Pharmacol. Exp. Ther.* 265: 1413-1420.
- Frühgott, R. E. and Jorhanian, D. (1983) Relation of cyclic GMP levels to endothelium-dependent relaxation by acetylcholine. *Fed. Proc.* 42: 619.
- Furchgott, R. F. and Zawadzki, J. V. (1980) The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature* 288: 373-376.
- Gandolfo, G. C., Romertino, S., Grotzmann, C., Van Luitelaar, G., Coenen, A., Buford, J. N. and Lazdunski, M. (1989) K^+ channel openers decrease seizures in genetically epileptic rats. *Eur. J. Pharmacol.* 167: 181-183.
- Garcla, G., Malta, A. and Seubert, R. (1990) European Patent Application. EP370 921A.
- Garland, C. J. and McPherson, G. A. (1992) Evidence that nitric oxide does not mediate the hyperpolarization and relaxation to acetylcholine in the rat small mesenteric artery. *Br. J. Pharmacol.* 105: 429-435.
- Garland, C. J., Plante, E., Kemp, B. K. and Cocks, T. M. (1995) Endothelium-dependent hyperpolarization: a role in the control of vascular tone. *Trends Pharmacol. Sci.* 16: 23-30.
- Gebremedhin, D., Ma, Y. A., Falck, J. R., Roman, R. J., Vanrolleghhe, M. and Harder, D. R. (1992) Mechanism of action of cerebral poyoxocastriacetic acids on cerebral arterial smooth muscle. *Am. J. Physiol.* 263: H319-H325.
- Gehlert, D. R. and Gackstein, S. L. (1991) Comparison of the distribution of binding sites for the potassium channel ligands [3H] apamin, [3H] charybdomycin and [3H] tetraethylammonium in the rat brain. *Neuroscience* 52: 191-205.
- Gelbard, C. H. and McCullough, J. B. (1992) Modulation of rabbit aortic Ca-activated K^+ channels by pinacidil, cromakalim and glibenclamide. *Am. J. Pharmacol.* 264: C1119-C1127.
- Ghaleb, B., Dabois-Randé, J. L., Hirtling, L., Giudicelli, J. F. and Bendaux, A. (1995) Comparisons of the effects of nicorandil, pinacidil, nicardipine and nitroglycerin on coronary vessels in the conscious dog: role of the endothelium. *Br. J. Pharmacol.* 114: 496-507.

- Glosson, M. M., Griffes, E. C., Nambi, P. and Ariyas, N. (1993) Endothelin stimulates ^{45}Ca efflux in rat glioma C6-Ba1 cells. *Neuropeptides* 26: 17-23.
- Gluck, Z., Winderman, F., Gutschow, M. P. and Wiedmann, P. (1985) Cardiovascular, endocrine and renal actions of acute potassium channel activation by BRL 14935 in man. *Cardiovasc. Drugs Ther.* 1: 241.
- Goddard, W. G. (1989) World Patent Application. WO/89/07571.
- Gopalakrishnan, M. and Triggle, D. J. (1990) The regulation of receptors, ion channels and proteins in congestive heart failure. *Cardiovasc. Drug Rev.* 8: 315-322.
- Gopalakrishnan, M., Rutledge, A., Kwon, Y. W., Butler, J., Fung, H. L., Janis, R. A. and Triggle, D. J. (1990) Regulation at the level of ATP-sensitive K^+ channels and 1,4-dihydropyridine-sensitive Ca^{2+} channels in experimental congestive heart failure secondary to myocardial infarction. *Pharmacologist* 12: 145.
- Gopalakrishnan, M., Janis, R. A. and Triggle, D. J. (1991) ATP-sensitive K^+ channels: pharmacologic properties and therapeutic potential. *Drug Dev. Res.* 28: 95-127.
- Gross, G. C., Duetting, G. J., Kemp, R. E. and Martin, T. J. (1987) Endothelin and the vasodilator action of calcitonin gene-related peptide (CGRP). *Br. J. Pharmacol.* 91: 729-733.
- Grofe, P., Quasthoff, S., Strupp, M. and Lehmann-Horn, F. (1990) Enhancement of K^+ conductance improves in vitro the contractility of skeletal muscle in hypokalemic periodic paralysis. *Muscle Nerve* 13: 451-457.
- Gross, T. L., Olmosky, C. J. and Howe, B. B. (1994) Anilide tertiary carbonyls: a novel series of K^+ channel openers. *Trends Pharmacol. Sci.* 15: 401-404.
- Greene, K. A., Foster, R. W. and Small, R. C. (1991) A patch clamp study of K^+ channel activity in bovine isolated tracheal smooth muscle cells. *Br. J. Pharmacol.* 102: 871-878.
- Greenwood, I. A. and Weston, A. H. (1993) Effects of rubidium ion responses to potassium channel openers in rat isolated aorta. *Br. J. Pharmacol.* 109: 925-932.
- Gross, E. (1977) Drugs acting on arterial smooth muscle (vasodilator drugs). In: *Antihypertensive Agents*, pp. 367-476. Gross, E. (ed.) Springer, Heidelberg.
- Gross, G. J. and Auchampach, J. A. (1992) Blockade of ATP-sensitive potassium channels prevents myocardial preconditioning in dogs. *Circ. Res.* 70: 221-231.
- Grever, G. J. (1994a) Promotive effects of ATP-sensitive potassium channel openers in experimental myocardial ischemia. *J. Cardiovasc. Pharmacol.* 24: S18-S27.
- Grever, G. J. (1994b) Promotive effects of ATP-sensitive potassium channel openers in models of myocardial ischemia. *Cardiovasc. Res.* 28: 773-782.
- Grever, G. J., McCulloch, J. R., Pfaller, A. J., Sargent, C. A. and Ardal, K. S. (1995) Cardioprotective profile of the cardiac-selective ATP-sensitive potassium channel opener BMS-180448. *J. Cardiovasc. Pharmacol.* 25: 469-50.
- Hamill, O. P., Marty, A., Sakmann, B. and Sigworth, F. J. (1981) Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflügers Arch.* 393: 85-100.
- Hamsters, T. C. and Weston, A. H. (1989) Cromakalim, nicotinic and pinacidil: novel drugs which open potassium channels in smooth muscle. *Gen. Pharmacol.* 20: 1-9.
- Haralson, T. C., Wei, S. W. and Weston, A. H. (1986) Comparison of the effects of BRL 14945 and verapamil on electrical and mechanical activity in rat portal vein. *Br. J. Pharmacol.* 88: 101-111.
- Harrison, C. S., Lutz, D. A. and Ducote, J. (1993) Potassium channel openers stimulate DNA synthesis in mouse epidermal keratinocytes and whole hair follicle cultures. *Skin Pharmacol.* 6: 170-178.
- Hasegawa, K., Redman, D. M., O'Brien, R. E. and McMurtry, I. F. (1990) Endothelin-1 causes pulmonary vasodilation in rats. *Am. J. Physiol.* 259: H48-H54.
- Hatton, R., Hays, C., Todd, M. H., Downing, O. A. and Whorl, K. A. (1993) Cromakalim reverses ischemic depression of skeletal muscle contractions in the rat hindlimb. *Br. J. Pharmacol. Proc. Suppl.* 102: 263P.
- Heaton, R. C., Weay, S. and Easter, D. A. (1993) Effects of metabolic inhibition and changes of intracellular pH on potassium permeability and contraction of rat uterus. *J. Physiol.* 465: 43-56.
- Heurteaux, C., Berridge, V., Widmann, C. and Lazdunski, M. (1991) K^+ channel openers prevent global ischemia-induced expression of c-fos, c-jun, heat shock protein and amyloid beta-protein precursor genes and neuronal death in rat hippocampus. *Proc. Natl. Acad. Sci. USA* 90: 9431-9435.
- Hockey, M., Bara, A. T., Bauersachs, J. and Busse, R. (1994) Characterization of EDHF as a cyclooxygenase/1450-derived arachidonic metabolite in mammals. *J. Physiol. (Lond)* 481: 407-414.
- Hedlund, H., Mariasson, A. and Andersson, K.-E. (1991) Effects of pinacidil on detrusor instability in men with bladder outlet obstruction. *J. Urol.* 146: 1345-1347.
- Hehl, S. and Neumcke, B. (1994) Diverse effects of pinacidil on K^+ channels in mouse skeletal muscle in the presence of different nucleotides. *Cardiovasc. Res.* 28: 841-846.
- Hille, B. (1994) *Ion Channels of Excitable Membranes*. Sinauer Associates, Sunderland.
- Hilton, S. M. (1959) A peripheral arterial constrictor mechanism underlying dilation of the femoral artery and concerned in functional dilation in skeletal muscle. *J. Physiol. (Lond)* 149: 93-111.
- Ho, K., Nichols, C. G., Lederer, W. J., Lytton, J., Vassilev, P. M., Kanaadskan, M. V. and Hebert, S. C. (1993) Cloning and expression of an inwardly-rectifying ATP-regulated potassium channel. *Nature* 363: 31-38.
- Ho, P. K., Towner, R. D., Eisenman, M. and Bernick, B. (1990) Pinacidil N^6 -cyclic- N^4 -pyridine- N^1 - N^2 -trimethylpropyl guanidine is a potent bronchodilator. *Eur. J. Pharmacol.* 183: 211-213.
- Hollingsworth, M., Amodeo, T., Edwards, G., Mironneau, J., Saito, J., P. Small, R. C. and Weston, A. H. (1987) The relaxation of BRL 14915 in rat uterus. *Br. J. Pharmacol.* 91: 803-813.
- Holmquist, E., Andersson, K.-E., Forsman, M. and Hedlund, H. (1988) K^+ channel openers for relaxation of isolated penile erectile tissue from rabbit. *J. Urol.* 144: 146-151.
- Holmquist, E., Andersson, K.-E. and Hedlund, H. (1990a) Effects of pinacidil on isolated human corpus cavernosus penis. *Acta Physiol. Scand.* 138: 461-469.
- Holmquist, E. (1988) Cyclic GMP is possible mediator of coronary arterial relaxation by nicotinic (SG-75). *J. Cardiovasc. Pharmacol.* 3: 364-370.
- Hong, K. W., Park, M. G., Shih, Y. W., Rham, B. Y. and Kuo, K. H. (1993) Effect of ouabain on relaxation induced by cromakalim on human and canine mesenteric arteries. *Eur. J. Pharmacol.* 231: 1-6.
- Howlett, D. B. and Longman, S. D. (1993) Identification of a binding site for 1-[4-chromakalim] in vascular and nonvascular smooth muscle cells. *Br. J. Pharmacol. Proc. Suppl.* 107: 96P.
- Hu, S., Kim, H. S. and Jeong, A. Y. (1991) Dual action of cromakalim on the Ca^{2+} -activated K^+ channel in smooth muscle cells of porcine coronary artery. *Eur. J. Pharmacol.* 194: 31-40.

- Huill, S. S., Kaiser, L., Jaffe, M. D. and Sparks, H. V. (1986) Endothelium-dependent flow-induced dilation of canine femoral and saphenous arteries. *Blood Vessels* 33: 183-198.
- Ichinose, M. and Barnes, P. J. (1990) A potassium channel activator modulates both excitatory noncholinergic and cholinergic neurotransmission in guinea-pig airways. *J. Pharmacol. Exp. Ther.* 252: 1207-1212.
- Jorda, M., Dewas, D. and McCulloch, J. (1991) Selective reduction of [125 I]pantoin binding sites in Alzheimer hippocampus: a quantitative autoradiographic study. *Brain Res.* 567: 51-56.
- Isoue, I., Nakaya, S. and Nakaya, Y. (1990) An ATP-sensitive K⁺ channel activated by extracellular Ca²⁺ and Mg²⁺ in primary cultured arterial smooth muscle cells. *J. Physiol. (Lond.)* 430: 132P.
- Ishikawa, J., Ichinose, M., Nakajima, N., Takahashi, T., Yamauchi, H., Miura, M. and Shirao, K. (1994) Potassium channel opener, YM-934, inhibits neurogenic plasma leakage in guinea-pig airways. *Am. J. Resp. Crit. Care Med.* 150: 1379-1383.
- Ito, S., Kajikuri, J., Ho, T. and Kuriyama, H. (1991) Effects of imakalim on changes in Ca²⁺ concentration and mechanical activity induced by noradrenaline in the rabbit mesenteric artery. *Br. J. Pharmacol.* 104: 227-233.
- Jungue, D., West, G. A., Gordan, E. L. and Winn, H. R. (1993) ATP-sensitive K⁺ channels in rat aorta and brain microvascular endothelial cells. *Am. J. Physiol.* 265: C812-C821.
- Jones, T. R., Charrier, L., Garcia, M. L. and Kaczorowski, G. J. (1990) Selective inhibition of relaxation of guinea-pig trachea by charybdotoxin a potent Ca²⁺-activated K⁺ channel inhibitor. *J. Pharmacol. Exp. Ther.* 255: 697-706.
- Kantor, P. F., Coetzee, W. A., Carmeliet, E. E., Denna, S. C. and Opie, L. H. (1993) Coupling of ATP-sensitive K⁺ channels to A₂ receptors by G-proteins in rat ventricular myocytes. *Circ. Res.* 66: 479-485.
- Kidney, J. C., Fuller, R. W., Worsell, Y. M., Lavender, E. A., Chung, K. F. and Barnes, P. J. (1993) Effect of an oral potassium channel activator, BRL 38227, on airway function and responsiveness in asthmatic patients - comparison with oral salbutamol. *Thorax* 48: 110-113.
- Kinoshita, M. and Sakai, K. (1990) Pharmacology and therapeutic effects of molsidomine. *Cardiovasc. Drugs* 4: 1075-1088.
- Kitama, K. and Kanouchi, M. (1993) K⁺ channel openers activate different K⁺ channels in vascular smooth muscle cells. *Cardiovasc. Drug Ther.* 7: 539-546.
- Klaus, E., Englert, H. C., Hoppe, M., Manis, D. and Zwergel, U. (1990) K⁺ channel openers inhibit the KCl-induced phasic rhythmic contractions in the upper urinary tract. *Eur. J. Pharmacol.* 183: 673-674.
- Robb, H. A. (1990) Potassium channels in excitable and non-excitable cells. *Rev. Physiol. Biochem. Pharmacol.* 115: 51-91.
- Komersova, K., Rogerson, J. W., Conway, E. L., Lim, T. C., Brown, D. J., Krum, H., Jackson, G. P., Murdoch, R. and Lous, W. J. (1995) The effects of levomecizolam (BRL 38227) on bladder function in patients with high spinal-cord lesion. *Br. J. Clin. Pharmacol.* 39: 207-209.
- Kume, H., Takai, A., Tokuno, H. and Tomita, T. (1989) Regulation of Ca²⁺ dependent K⁺ channel activity in tracheal myocytes by phosphorylation. *Nature* 341: 152-154.
- Kuriuchi, Y., Nakajima, T. and Sugimoto, T. (1986) On the mechanism of activation of muscarinic K⁺ channel by adenosine in isolated smooth cells - involvement of GTP-binding proteins. *Phylog. Arch.* 407: 264-274.
- Kuriyama, H. and Suzuki, H. (1978) The effects of acetylcholine on the membrane and contractile properties of smooth muscle cells of the rabbit superior mesenteric artery. *Br. J. Pharmacol.* 65: 493-501.
- Laurens, F., Michel, A., Bonnet, P. A., Chaput, J. P. and Boucard, M. (1993) Evaluation of the relaxant effects of SCA40, a novel charybdotoxin-sensitive potassium channel opener, in guinea-pig isolated trachealis. *Br. J. Pharmacol.* 108: 622-626.
- Lawson, K. and Castero, I. (1989) Effects of Ca²⁺ antagonists and K⁺ channel activators on K⁺-induced contractions of the rat aorta. *J. Auton. Pharmacol.* 9: 329-336.
- Lawson, K. and Hicks, P. E. (1993) Potassium channel openers: pharmacological anomalies suggest heterogeneous sites of action. *Curr. Opin. Invest. Drugs* 2: 1209-1216.
- Lawson, K., Barras, M., Zaid-Sudic, E., Martin, D. J., Armstrong, J. M. and Hicks, P. E. (1992) Differential effects of endothelin on the vasorelaxant properties of benzopyran and non-benzopyran potassium channel openers. *Br. J. Pharmacol.* 107: 58-65.
- Lawson, K., Barras, M., Armstrong, J. M. and Hicks, P. E. (1993a) Endothelin-dependent vasorelaxation to BRL 38227 in rat aorta contracted with prostaglandin F_{2α}. *Fundam. Clin. Pharmacol.* 7: 183.
- Lawson, K., Barras, M., Armstrong, J. M. and Hicks, P. E. (1993b) Effects of glibenclamide and charybdotoxin on relaxation to the potassium channel activator BRL 55834 in guinea-pig trachea. *Br. J. Pharmacol. Proc. Suppl.* 110: 237.
- Lawson, K., Barras, M., Armstrong, J. M. and Hicks, P. E. (1995) Effects of NS-004, an activator of BK_{Ca} channels, in guinea-pig isolated trachea. *Fundam. Clin. Pharmacol.*, in press.
- Lebel, M., Grosse, J. H. and Lacourcière, Y. (1989) Short term monotherapy with the potassium channel activator BRL 34915 on endocrine sodium regulation in essential hypertension. *Am. J. Hypertens.* 2: 256-258.
- Lei, Y. H., Barnes, P. J. and Rogers, D. E. (1993) Inhibition of neurogenic plasma exudation and bronchoconstriction by a K⁺ channel activator, BRL 38227, in guinea-pig airways *in vivo*. *Eur. J. Pharmacol.* 239: 257-259.
- Lijnen, P., Fagard, R., Staessen, J., Weirich, T., Moerman, E. and Amery, A. (1989) Humoral and cellular effects of the K⁺ channel activator cromakalim in man. *Eur. J. Clin. Pharmacol.* 37: 609-611.
- Lippton, H. L., Cohen, G. A., McMurtry, J. F. and Hyman, A. L. (1991) Pulmonary vasodilation to endothelin isopeptides is seen in mediated by potassium channel activation. *J. Appl. Physiol.* 70: 947-952.
- Longman, S. D. and Hamblin, T. C. (1992) Potassium channel activator drugs: mechanism of action, pharmacological properties and therapeutic potential. *Med. Res. Rev.* 12: 73-148.
- Luckhoff, A. and Bause, R. (1990) Activators of potassium channels enhance calcium influx into endothelial cells as a consequence of potassium currents. *Naunyn-Schmiedeberg Arch. Pharmacol.* 342: 94-99.
- MacMillan, S., Sheridan, R. D., Chilvers, E. R. and Patmore, L. (1995) A comparison of the effects of SCA40, NS 004 and NS 1619 on large conductance Ca²⁺-activated K⁺ channels in bovine tracheal smooth muscle cells in culture. *Br. J. Pharmacol.* 116: 1656-1660.
- Malmgren, A., Andersson, K.-E., Sjogren, C. and Andersson, P. O. (1989) Effects of yohimbine and cromakalim (BRL 34915) on bladder function in rats with detrusor instability. *J. Urol.* 141: 1134-1138.

- Masuda, Y., Yoshitomi, M., Ishimura, Y., Karch, I. and Oka, M. (1994) Effects of the potassium channel openers cromakalim and pinacidil on catecholamine secretion and calcium mobilization in cultured bovine adrenal chromaffin cells. *Biochem. Pharmacol.* 47: 1751-1758.
- Matsu, D., Nagata, T., Rogawski, M. A. and Avoli, M. (1994) Potassium channel activators counteract anoxic hyperexcitability but not 4-aminopyridine induced epileptiform activity in the rat hippocampal slice. *Neuropharmacology* 33: 1515-1522.
- McCaig, J. D. and Welsh, M. J. (1986) Calcium-activated potassium channels in canine smooth muscle. *J. Physiol.* 372: 113-127.
- McKay, M. C., Dzeravsky, S. I., Meanwell, N. A., Olesen, S.-P., Reinhardt, P. H., Levitan, I. B., Adelman, J. P. and Grinvald, V. K. (1994) Opening of large-conductance calcium-activated potassium channels by the substituted benzimidazole, NS-004. *J. Neurophysiol.* 71: 1823-1882.
- McLean, M. J., Schmutz, M., Wami, A. W., Olpe, H. R., Fortes, C. and Feldmann, K. F. (1994) Oxcarbazepine - mechanism of action. *Epilepsia* 35: 55-59.
- McManus, O. B., Harris, G. H., Gangiacomo, K. M., Egerdshaus, P., Reuben, J. P., Adity, M. E., Durka, J. E., Kaczorowski, G. J. and Garcia, M. L. (1993) An activator of calcium-dependent potassium channels isolated from a medicinal herb. *Biochemistry* 32: 6128-6133.
- Meers, P., Arner, K., Sanbous, B. and Toso, L. (1994) Activation of maxi-K channel of human myometrial cells by the hormone relaxin. *Physiol. J.* 66: A207.
- Mellin, C. and Wilson, W. S. (1991) Comparison of the effects of vasodilator drugs on intracranial pressure and vascular relaxation. *Br. J. Pharmacol. Clin. Suppl.* 104: 55P.
- Müller, B. J. (1990) Glucose-regulated potassium channels are sweet morsels for neurobiologists. *Trends Neurosci.* 13: 197-199.
- Mura, M., Belvisi, M. G., Stretton, D., Yacoub, M. H. and Barnes, P. J. (1991) Role of potassium channels in bronchodilator responses in human airways. *Am. Rev. Resp. Dis.* 146: 132-136.
- Miyoshi, Y., Nakaya, Y., Wakatsuki, T., Nakaya, S., Fujino, K., Sano, K. and Inoue, I. (1992) Endothelin blocks ATP-sensitive K⁺ channels and depolarizes smooth muscle cells of porcine coronary artery. *Circ. Res.* 70: 612-616.
- Monga, M. and Gross, R. K. (1993) Management of preeclampsia. *Semin. Perinatol.* 19: 84-96.
- Morley, J. (1994) K⁺ channel openers and suppression of airway hyperactivity. *Trends Pharmacol. Sci.* 15: 461-468.
- Morley, J. E. (1980) The neuroendocrine control of appetite: the role of the endogenous opiates, cholecystikinin, TRH, gamma-aminobutyric acid and the diuretic receptor. *Life Sci.* 27: 355-368.
- Morrison, J. J., Ashford, M. L., Khan, R. N. and Smith, S. K. (1993) Effects of PGOs on isolated pregnant human myometrium before and after the onset of labor: potential of oxytocin. *Am. J. Obstet. Gynecol.* 169: 1277-1285.
- Mutter, A. M. and Würel, M. (1983) Mode of action of cycloheximide and triamterene. Ex vivo effect on Na⁺ and Rb⁺ efflux from periental smooth muscle. *Eur. J. Pharmacol.* 66: 129-133.
- Müller-Bühl, U., Diehm, C., Kraus, T., Zimmermann, R., Moes, H. and Exner, H. H. (1985) Clinical effects of isoproterenol in patients with intracranial obstruction. *Eur. J. Clin. Pharmacol.* 28: 127-131.
- Murray, M. A., Ficker, R. W. and Small, R. C. (1990) Effects of the K⁺ channel openers cromakalim and RP 49356 on the cyclic nucleotide content of guinea-pig isolated trachealis muscle. *Br. J. Pharmacol.* 100: 161P.
- Nakajima, S., Kurokawa, K., Imamura, N. and Ueda, M. (1989) A study on the hypotensive mechanism of pinacidil - relationship between its vasodilating effect and intracellular Ca²⁺ levels. *Jpn. J. Pharmacol.* 49: 205-211.
- Nakashima, M. and Vanhoutte, P. M. (1993a) Endothelin-1 and endothelin-1 cause endothelin-dependent hyperpolarization in the rat mesenteric artery. *Am. J. Physiol.* 265: H2137-H2141.
- Nakashima, M. and Vanhoutte, P. M. (1993b) Endothelin and membrane potential of vascular smooth muscle of the canine coronary artery. *J. Cardiovasc. Pharmacol.* 22: S229-S233.
- Nelson, M. T. (1993) Ca²⁺-activated potassium channels and ATP-sensitive potassium channels as modulators of vascular tone. *Trends Cardiovasc. Med.* 3: 54-60.
- Nelson, M. T., Huang, Y., Brayden, J. E., Hescheler, J. and Strassman, N. B. (1990) Activation of K⁺ channels is involved in arterial dilations to calcitonin gene-related peptide. *Nature* 344: 770-771.
- Newgreen, D. T., Brag, K. M., McHarg, A. D., Weston, A. H., Durr, S., Brown, B. S., Kay, R. B., Edwards, C., Longmore, J. and Southerton, J. S. (1990) The actions of diltiazem and mexiletine sulphate on rat blood vessels: a comparison with cromakalim. *Br. J. Pharmacol.* 100: 605-613.
- Nielsen-Kudsk, J., Bang, L. and Brønsgaard, A. M. (1990) Glibenclamide blocks the relaxant action of pinacidil and cromakalim in airway smooth muscle. *Eur. J. Pharmacol.* 180: 291-296.
- Nielsen-Kudsk, J. E., Møller-Nielsen, S. and Thirstrup, S. (1994) Inhibition by cromakalim, carbutaline, theophylline and verapamil of noncholinergic nerve-mediated contractions of guinea-pig isolated bronchi. *Pulm. Pharmacol.* 7: 285-292.
- Noma, A. (1983) ATP-regulated K channels in cardiac muscle. *Nature* 305: 147-148.
- North, R. A. (1989) Drug receptors and the inhibition of nerve cells. *Br. J. Pharmacol.* 98: 11-28.
- Oates, J. A., Connolly, M. E., Pritchard, B. N. C., Shand, D. G. and Schapel, G. (1977) The clinical pharmacology of antihypertensive drugs. In: *Antihypertensive Agents*, pp. 571-632. Cross, F. (ed.) Springer, Heidelberg.
- Okenstedt, H., Kraus, T., Schuler, M. and Böcher, G. (1989) Clinical benefits of isoproterenol, a stable prostacyclin (PGI₂) analog, in severe peripheral arterial disease (PAD). *Adv. Prostaglandin Thromboxane Leukot.* 10: 311-316.
- Oscar, M., Posa, E. D., Barros, M., Robles, L. I. and Boeyens, J. M. (1990) An ATP-dependent potassium channel blocker antagonizes morphine analgesia. *Eur. J. Physiol.* 156: 377-378.
- Oscar, M., Del Pozo, E. and Boeyens, J. M. (1989) ATP-dependent K channel blockers antagonize morphine but not U-50488B-induced antinociception. *Eur. J. Pharmacol.* 210: 203-207.
- Quae, K., Kujala, S., Nakao, K., Kurokawa, K., Kurivama, H. and Weston, A. H. (1990) Action of cromakalim on ionic currents recorded from single smooth muscle cells of the rat portal vein. *J. Pharmacol. Exp. Ther.* 252: 812-819.
- Ozkan, Y., Yangrawata, T. and Tama, N. (1993) BRL 28277 (kainate)-induced hyperpolarization reduces the sensitivity to Ca²⁺ contractile elements in canine coronary artery. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 347: 438-444.
- Olesen, S.-P. and Wajsb, F. (1992) European Patent Application E. P0477819A2.
- Olesen, S.-P., Wajsb, F. and Hayes, A. G. (1993) A novel benzimidazole, NS004, activates large-conductance Ca²⁺-dependent K channels in aortic smooth muscle. *Br. J. Pharmacol. Proc. Suppl.* 110: 23P.

- Olesen, S.-P., Munch, E., Wassen, F. and Drejer, J. (1994a) NS-004 - an activator of Ca^{2+} dependent K channels in cerebellar granule cells. *Neuroreport* 5: 1001-1004.
- Olesen, S.-P., Munch, E., Wassen, F. and Drejer, J. (1994b) Selective activation of Ca^{2+} dependent K channels by novel benzimidazole. *Eur. J. Pharmacol.* 251: 55-59.
- Parlocke, R. M., Spencer, A. M., Gater, P. R. and Waterfall, J. F. (1992) PAF-induced hyperactivity in the guinea-pig is inhibited by Ro 31-6930 and salmeterol. In: *New Concepts in Asthma*, Vol. 5, Vargafik, B. B., Tarayre, J. P. and Tisue-Versaille, J. (eds.) MacMillan, London.
- Palmer, R. M. J., Feenys, A. G. and Morcadas, S. (1987) Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature* 327: 524-526.
- Patelunas-Hoffman, D. M., Carmist, W. J., Colatsky, T. J. and Perinich, R. L. (1994) Analysis of the potassium channel opensers celastrol, pinacidil and cromakalim in platelet models of thrombosis. *Thromb. Res.* 74: 441-452.
- Passi, T., Kaukinen, S., Ylitalo, P., Reinikainen, P. and Vapalahti, H. (1986) Supro, a prostaglandin analog in the treatment of advanced peripheral arterial disease. *Vasc. Surg.* 20: 301-306.
- Pickkers, P. and Hughes, A. D. (1995) Relaxation and decrease in $[\text{Ca}^{2+}]_i$ by hydrochlorothiazide in guinea-pig isolated mesenteric arteries. *Br. J. Pharmacol.* 114: 703-707.
- Piper, L., Minshall, E., Downing, S. J., Hollingsworth, M. and Sadrieh, H. (1993) Effects of several potassium channel openers and glibenclamide on the uterus of the rat. *Br. J. Pharmacol.* 101: 901-907.
- Piper, L., Downing, S. J. and Hollingsworth, M. (1992) Cross tolerance between cromakalim and RP 49356 in the uterus of the rat in vivo and in vitro. *Eur. J. Pharmacol.* 219: 347-353.
- Pohl, V., Heitz, J., Busse, R. and Zauscher, E. (1986) Crucial role of the endothelium in the vasodilator response to increased flow in vivo. *Hypertension* 8: 37-44.
- Popoli, P., Perzola, A., Sagratella, S., Zeng, Y. C. and Scotti De Carolis, A. (1991) Cromakalim (BRL 34915) counteracts the epiphriform activity elicited by debrisoquine and verapamil in rats. *Br. J. Pharmacol.* 104: 907-913.
- Poucelier, M. (1993) *European Patent Application E. P. 370902*.
- Quast, U. and Baumlin, Y. (1988) Comparison of the effects of 9-K^+ and M^{Rb^+} elicited by cromakalim (BRL 34915) in tonic and phasic vascular tissue. *Naunyn-Schmiedeberg Arch. Pharmacol.* 338: 319-326.
- Quast, U., Bray, K. M., Baumlin, Y. and Dosogne, J. (1992) Potassium channel openers: pharmacology and therapeutic prospects. In: *Trends Receptor Research*, pp. 309-332, Angeli, E., Guilleini, U. and Quaglia, W. (eds.) Elsevier Science Publishers B.V., Amsterdam.
- Quast, U., Bray, K. M., Andres, H., Manley, P. W., Baumlin, Y. and Dosogne, J. (1993) Binding of the K^+ channel opener [14] P1075 in rat isolated aorta: relationship to functional effects of openers and blockers. *Mol. Pharmacol.* 43: 474-481.
- Quasthoff, S., Spuler, A., Lehmann-Horn, F. and Grafe, P. (1989) Cromakalim, pinacidil and RP 49356 activate a tetraethylammonium-sensitive K^+ conductance in human skeletal muscle fibres. *Pflügers Arch.* 414: S179-S180.
- Quasthoff, S., Spuler, A., Spiermeister, W., Lehmann-Horn, F. and Grafe, P. (1990) K^+ channel openers suppress myotonic activity of human skeletal muscle in vitro. *Eur. J. Pharmacol.* 186: 125-128.
- Rabjohn, D. and Karlsson, J. A. (1991) Potassium channel openers: airway pharmacology and clinical possibilities in asthma. *Prog. Drug Res.* 37: 161-180.
- Ramalli, M. D. (1991) Vascular activities of the endothelins. *Pharmacol. Ther.* 50: 73-93.
- Randall, M. D. and Griffith, T. M. (1993) Modulation of vasodilation to levcromakalim by hypoxia and EDRF in the isolated perfused rabbit ear: a comparison with pinacidil, sodium nitroprusside and verapamil. *Br. J. Pharmacol.* 109: 336-339.
- Rapaport, R. M. and Murad, F. (1983) Agonist-induced endothelium-dependent relaxation in rat thoracic aorta may be mediated through cGMP. *Circ. Res.* 52: 352-357.
- Renne, H. L., Vaughan, R. F. and Peers, C. (1992) Glibenclamide inhibits voltage-gated K^+ current in the human neuroblastoma cell line SH-SY5Y. *Neurosci. Lett.* 135: 37-40.
- Renard, J. F., Desmuelle, C., Schmid-Aumarchi, H., Hugues, M., Serrurier, G. and Lazdunski, M. (1986) Expression of a myotonic receptor in muscles of patients with myotonic muscle dystrophy. *Nature* 319: 678-680.
- Richen, C., Pratz, J., Mulder, R., Mondot, S., Giudicelli, J. F. and Caverio, I. (1990) Cardiovascular and biological effects of K^+ channel openers: a class of drugs with vasorelaxant and cardio-protective properties. *Life Sci.* 47: 1693-1705.
- Robertson, D. W. and Steinberg, M. L. (1994) Potassium channel modulators: scientific applications and therapeutic promise. *J. Med. Chem.* 37: 1529-1541.
- Rudy, B. (1988) Diversity and ubiquity of K channels. *Neuroscience* 25: 729-749.
- Sablayrolles, C., Bonnet, P. A., Cros, G., Chupat, J.-P. and Boucard, M. (1988) *World Patent Application WO 88/04298*.
- Sadmei, H. and Beech, D. J. (1995) *low currents and inhibitory effects of glibenclamide in animal vesicle smooth muscle cells*. *Br. J. Pharmacol.* 115: 1447-1454.
- Sanborn, B. M. (1995) Ion channels and the control of myocardial electrical activity. *Sem. Perinatol.* 19: 31-40.
- Santucci, P. and Maggi, C. A. (1994) Inhibitory transmitter action of calcitonin gene related peptide in guinea-pig ureter via activation of glibenclamide-sensitive K^+ channels. *Br. J. Pharmacol.* 113: 588-592.
- Sargent, C. A., Grover, G. J., Antonacci, M. J. and McCullough, J. R. (1993) The cardioprotective, vasorelaxant and electrophysiological profile of the large-conductance calcium-activated potassium channel opener NS-004. *J. Pharmacol. Exp. Ther.* 266: 1422-1429.
- Savini, J.-P. and Marthan, R. (1993) Effect of cromakalim on KCl , noradrenaline- and angiotensin II-induced contractions in the rat pulmonary artery. *Pulm. Pharmacol.* 6: 41-48.
- Schmid-Aumarchi, H., Amoroso, S., Fasser, M. and Lazdunski, M. (1990) K^+ channel openers activate brain sulfonylurea-sensitive K^+ channels and block neurotransmitter. *Proc. Natl. Acad. Sci. USA* 87: 3489-3492.
- Sellers, A. J. and Ashford, M. L. J. (1994) Activation of BK_{Ca} channels in acutely dissociated neurones from the rat ventromedial hypothalamus by NS 1619. *Br. J. Pharmacol.* 113: 659-661.
- Sellers, A. J., Boden, P. R. and Ashford, M. L. J. (1992) Lack of effect of potassium channel openers on ATP-modulated potassium channels recorded from rat ventromedial hypothalamic neurones. *Br. J. Pharmacol.* 107: 1068-1074.
- Shah, A. M. (1992) Vascular endothelium. *Br. J. Hosp. Med.* 48: 540-549.
- Sheppard, D. N. and Welsh, M. J. (1992) Effects of ATP-sensitive K^+ channel regulators on cyclic nucleotide transmembrane covalence regulator chloride currents. *J. Gen. Physiol.* 100: 573-591.

- Stiegel, G., Mironneau, J., Scholke, F., Lomand, G. and Stock, G. (1990) Potassium channel opening and vasorelaxation. In: *Endothelium-Derived Relaxing Factors*, pp. 265-280. Raharjy, G. M. and Vanhoutte, E. P. M. (eds) Karger, Basel.
- Singer, D. R. J., Markand, N. D., Miller, M. A., Sugden, A. L. and MacGregor, G. A. (1989) Potassium channel stimulation in normal subjects and in patients with essential hypertension: an acute study with cromakalim (BRL 34915). *J. Hypertens.* 7: 5294-5295.
- Small, R. C., Berry, J. L., Barka, J. F., Cook, S. J., Foster, R. W., Green, K. A. and Murray, A. A. (1992) Potassium channel activators and bronchial asthma. *Clin. Exp. Allergy* 22: 11-18.
- Smith, H. (1989) Animal models of asthma. *Pain. Pharmacol.* 2: 59-74.
- Smith, H. (1992) Asthma, inflammation, eosinophils and bronchial hyperresponsiveness. *Clin. Exp. Allergy* 22: 187-197.
- Southerton, J. S., Weston, A. H., Bray, K. M., Newgreen, D. T. and Taylor, S. G. (1988) The potassium channel opening action of pinacidil: studies using biochemical, ion flux and microelectrode techniques. *Naunyn-Schmiedeberg Arch. Pharmacol.* 338: 310-318.
- Spuler, A., Lehmann-Horn, F. and Gräbe, P. (1989) Cromakalim (BRL 34915) restores in vitro the membrane potential of depolarized human skeletal muscle fibres. *Naunyn-Schmiedeberg Arch. Pharmacol.* 339: 327-331.
- Standen, N. B., Quayle, J. M., Davies, N. W., Brayden, J. E., Huang, Y. and Nelson, M. T. (1989) Hyperpolarizing vasodilators activate ATP-sensitive K⁺ channels in arterial smooth muscle. *Science* 249: 377-380.
- Tate, M., Parkinson, H. C., Coleman, H. A., Nickl, T. O. and Dunning, G. J. (1990) Hyperpolarization and relaxation of arterial smooth muscle is caused by nitric oxide derived from the endothelium. *Nature* 346: 69-71.
- Taylor, S. G. and Weston, A. H. (1988) Endothelium-derived hyperpolarizing factor: a new endogenous inhibitor from the vascular endothelium. *Trends Pharmacol. Sci.* 9: 272-274.
- Taylor, S. G., Arty, J. R. S., Bond, J., Buckle, D. R., Shaw, D. J., Taylor, J. F. and Ward, J. (1992) The inhibitory effects of cromakalim and its active congener BRL 38227 against various spasmogens in guinea-pig and human airways: comparison with pinacidil and verapamil. *J. Pharmacol. Exp. Ther.* 261: 429-437.
- Templeton, A. G. B., MacLennan, S. J., Fan, M. A., MacMillan, S., Chilvers, E. R. and Hicks, P. E. (1989) A NS1619, a BK_v channel opener in human bronchus. *Br. J. Pharmacol. Proc. Suppl.* 115: 119P.
- Thomas, P., Devion, M. S. and Sheridan, D. J. (1990) Acute haemodynamic effects of cromakalim in patients with angina pectoris. *Br. J. Pharmacol.* 105: 324-331.
- Tombas, C. F., Norman, N. R., Cropp, V. E., Lee, K. S., Gadowood, R. C. and Shebuts, R. C. (1992) Limitation of myocardial injury by the potassium channel opener cromakalim and the non-vascular analog BRL 34915: vascular endothelial actions in vitro and in vivo. *J. Pharmacol. Exp. Ther.* 261: 1261-1268.
- Towle, A., Sordahl, P. and Day, D. E. (1992) Reperfusion-induced arrhythmias and myocardial ischaemia: a pharmacological interaction between pinacidil and calcitonin in isolated rat hearts. *Basic Res. Cardiol.* 87: 365-364.
- Towle, A., Sordahl, P., Engelmann, R. M. and Day, D. E. (1993) Potassium channel openers and blockers: do they possess antiarrhythmic or antiarrhythmic activity in ischaemic and reperfusion rat hearts. *J. Pharmacol. Exp. Ther.* 267: 1355-1362.
- Trigo-Recha, F., Hsu, G. L., Donatucci, C. E., Martinez-Miranda, L., Lee, T. E. and Fingberg, E. A. (1990) Intracellular mechanism of penile erection in monkeys. *Neurosci. Urology* 14: 71-80.
- Tzifant, H. A., Malmert, W., Fleischacker, A. and Adelwahrer, N. (1991) Potassium channels and modulating factors of channel functions in the human myocardium. *Z. Kardiol.* 80: 29-31.
- Tslenko, T. N. and Bulecki, R. (1989) Altered potassium channel ion fluxes to vasoactive stimuli following: collateral enrichment in arterial smooth muscle. *Arteriosclerosis* 9: 769.
- Usune, S., Katsuragi, T. and Furukawa, T. (1991) Involvement of K⁺ channel opening in endothelin-1 induced suppression of spontaneous contractions in the guinea-pig taenia coli. *Can. J. Physiol. Pharmacol.* 69: 1908-1913.
- Vanden Burg, M. J., Woodward, S. P., Hassan, M., Stewart-Lung, P. M. and Tasker, T. C. G. (1986) Potassium channel activators lower blood pressure—in an initial study of BRL 34915 in hypertensive patients. *J. Hypertens.* 4: S166-S167.
- Vanden Burg, M. J., Woodward, S. P. and Stewart-Lung, P. M. (1987) Potassium channel activators antihypertensive activity and adverse effect profile of BRL 34915. *J. Hypertens.* 5: S193-S195.
- Van Reuterghem, C., Vigue, P., Barbonin, J., Schaud-Althaus, A., Fiehn, C. and Lardoux, M. (1988) Molecular mechanism of action of the vasoconstrictor peptide endothelin. *Biochem. Biophys. Res. Commun.* 157: 977-985.
- Vergani, A. V., Scamm, A. and Berman, A. (1992) Pinacidil potentiates morphine analgesia. *Life Sci.* 50: 119-138.
- Walden, D. J., Koweb, T. T., Barker, C. A., Johnson, G. A. and Buhl, A. E. (1993) Enhanced in vitro hair growth at the air liquid interface: pinacidil preserves the root sheath in cultured whisker follicles. *In Vitro Cell Dev. Biol.* 29A: 555-561.
- Ward, J. F. T., Taylor, S. G. and Collier, M. L. (1992) The novel benzopyranol potassium channel activator BRL 38227 activates two potassium channels in bovine aortic smooth muscle. *Br. J. Pharmacol. Proc. Suppl.* 167: 49P.
- Wareham, A. C. (1992) Skeletal muscle potassium channels and their relevance to muscle disease. In: *Potassium Channel Modulators: Pharmacological Molecular and Clinical Aspects*, pp. 213-271. Weston, A. H. and Hamilton, T. C. (eds) Blackwell Scientific, Oxford.
- Waucho, C. J., Dockrell, M. F. C., Haynes, W. G., Olverman, H. J., Williams, B. C. and Webb, D. J. (1992) The potassium channel opener BRL 38227 inhibits binding of [³H]-labelled endothelin-1 to rat cardiac membranes. *Biochem. Biophys. Res. Commun.* 185: 630-633.
- Wick, R. and Neumker, B. (1990) Effects of potassium channel openers on single channels in mouse skeletal muscle. *Naunyn-Schmiedeberg Arch. Pharmacol.* 347: 758-763.
- Wein, A. J. (1993) Practical pharmacology. *Lancet Clin. North Am.* 336: 289-291.
- Wein, A. H. and Weston, A. H. (1989) The effects of BRL 34915 and nicotinic acid on electrical and mechanical activity and on "Rb" efflux in rat blood vessels. *Br. J. Pharmacol.* 88: 121-128.
- Wickler, B. R., Ley, C. W. and Jaffe, E. A. (1978) Stimulation of endothelial cell protein production by theophylline, trypsin and the ionophore A23187. *J. Clin. Invest.* 62: 923-932.
- Wick, S. P. and Dunlow, L. D. (1993) Antinociceptive activity of intracutaneously administered potassium channel openers and opioid agonists: a common mechanism of action. *J. Pharmacol. Exp. Ther.* 267: 390-399.

- Weselcouch, E. O., Sargents, C., Wilde, M. W. and Smith, M. A. (1993) ATP-sensitive potassium channels and skeletal muscle function *in vitro*. *J. Pharmacol. Exp. Ther.* 267: 410-416.
- Wessler, L., Holtz, C., Mardigan, J., Pohar, D., Reinheimer, T. and Racke, K. (1993) Cromakalim inhibits electrically evoked [^{3}H] acetylcholine release from a tube preparation of the rat isolated trachea by an epithelium dependent mechanism. *Naunyn-Schmiedeberg Arch. Pharmacol.* 346: 14-20.
- Wickenden, A. D., Grimwood, S., Grant, I. L. and Todd, M. H. (1991) Comparison of the effects of the K^+ channel openers cromakalim and minoxidil sulphate on vascular smooth muscle. *Br. J. Pharmacol.* 103: 1148-1152.
- Wilde, A. A. M. (1994) Proarrhythmic potential of the potassium channel openers. *Cardiovasc. Res.* 28: 918-929.
- Williams, A. J., Lee, T. H., Cochrane, G. M., Hopkirk, A., Vyse, T., Chiew, F., Laverdes, E., Richards, D. H., Owen, S., Stone, P., Church, S. and Woodcock, A. A. (1990) Attenuation of nocturnal asthma by cromakalim. *Lancet* 336: 334-336.
- Wilson, C. (1989) Inhibition by sulphonylureas of vasorelaxation induced by K^+ channel activators *in vitro*. *J. Auton. Pharmacol.* 9: 71-78.
- Yamagishi, T., Yanagisawa, T. and Taira, N. (1992) K^+ channel openers, cromakalim and K4032, inhibit agonist-induced Ca release in canine coronary artery. *Naunyn-Schmiedeberg Arch. Pharmacol.* 346: 691-700.
- Yao, Z. H. and Gross, G. J. (1994) Effects of the K_{ATP} channel opener hirsakalim on coronary blood flow, monophasic action potential duration and infarct size in dogs. *Circulation* 89: 1769-1775.
- Zini, S., Relais, A. B., Arnergaard, C. and Ben-Ari, Y. (1993) Effect of potassium channel modulators on the release of glutamate induced by ischaemic-like conditions in rat hippocampal slices. *Neurosci. Lett.* 153: 202-205.
- Zona, C., Tinarelli, V., Palma, E., Pirroni, G. C. and Avoli, M. (1990) Potassium currents in rat cortical neurons: cultures are enhanced by the antiepileptic drug carbamazepine. *Can. J. Physiol. Pharmacol.* 68: 545-547.
- Zurkier, B. J., Lenzen, S., Mannen, K., Panten, U. and Trube, G. (1988) Concentration-dependent effects of tolbutamide, glibenclamide, glibenclamide and diazoxide on ATP-regulated K^+ currents in pancreatic β cells. *Naunyn-Schmiedeberg Arch. Pharmacol.* 337: 225-230.